

7. References

- Adams, D. et al., 2012. BLUEPRINT to decode the epigenetic signature written in blood. *Nature biotechnology*, 30(3), pp.224–226. Available at: <http://dx.doi.org/10.1038/nbt.2153>.
- Akira, S., Uematsu, S. & Takeuchi, O., 2006. Pathogen recognition and innate immunity. *Cell*, 124(4), pp.783–801. Available at: <http://dx.doi.org/10.1016/j.cell.2006.02.015>.
- Alipanahi, B. et al., 2015. Predicting the sequence specificities of DNA- and RNA-binding proteins by deep learning. *Nature biotechnology*, 33(8), pp.831–838. Available at: <http://dx.doi.org/10.1038/nbt.3300>.
- Anderson, C.A. et al., 2011. Synthetic associations are unlikely to account for many common disease genome-wide association signals. *PLoS biology*, 9(1), p.e1000580. Available at: <http://dx.doi.org/10.1371/journal.pbio.1000580>.
- Atabani, S.F. et al., 2005. Association of CTLA4 polymorphism with regulatory T cell frequency. *European journal of immunology*, 35(7), pp.2157–2162. Available at: <http://dx.doi.org/10.1002/eji.200526168>.
- Bannister, A.J. & Kouzarides, T., 2011. Regulation of chromatin by histone modifications. *Cell research*, 21(3), pp.381–395. Available at: <http://dx.doi.org/10.1038/cr.2011.22>.
- Barski, A. et al., 2007. High-Resolution Profiling of Histone Methylations in the Human Genome. *Cell*, 129(4), pp.823–837. Available at: <http://www.cell.com/article/S0092867407006009/abstract> [Accessed August 18, 2018].
- Benevolenskaya, E.V., 2007. Histone H3K4 demethylases are essential in development and differentiation. *Biochemistry and cell biology = Biochimie et biologie cellulaire*, 85(4), pp.435–443. Available at: <http://dx.doi.org/10.1139/O07-057>.
- Bernstein, B.E. et al., 2010. The NIH Roadmap Epigenomics Mapping Consortium. *Nature biotechnology*, 28(10), pp.1045–1048. Available at: <http://dx.doi.org/10.1038/nbt1010-1045>.
- Bini, D.A., Latouche, G. & Meini, B., 2005. INTRODUCTION TO MARKOV CHAINS. In *Numerical Methods for Structured Markov Chains*. Oxford: Oxford University Press. Available at:

<http://www.oxfordscholarship.com/10.1093/acprof:oso/9780198527688.001.0001/acprof-9780198527688-chapter-1>.

- Blattler, A. et al., 2014. Global loss of DNA methylation uncovers intronic enhancers in genes showing expression changes. *Genome biology*, 15(9), p.469. Available at: <http://dx.doi.org/10.1186/s13059-014-0469-0>.
- Bluestone, J.A., Tang, Q. & Sedwick, C.E., 2008. T regulatory cells in autoimmune diabetes: past challenges, future prospects. *Journal of clinical immunology*, 28(6), pp.677–684. Available at: <http://dx.doi.org/10.1007/s10875-008-9242-z>.
- Bock, C. & Lengauer, T., 2008. Computational epigenetics. *Bioinformatics*, 24(1), pp.1–10. Available at: <http://dx.doi.org/10.1093/bioinformatics/btm546>.
- Browning, S.R. & Browning, B.L., 2007. Rapid and accurate haplotype phasing and missing-data inference for whole-genome association studies by use of localized haplotype clustering. *American journal of human genetics*, 81(5), pp.1084–1097. Available at: <http://dx.doi.org/10.1086/521987>.
- Buenrostro, J.D. et al., 2013. Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. *Nature methods*, 10, p.1213. Available at: <http://dx.doi.org/10.1038/nmeth.2688>.
- Bush, W.S. & Moore, J.H., 2012. Chapter 11: Genome-wide association studies. *PLoS computational biology*, 8(12), p.e1002822. Available at: <http://dx.doi.org/10.1371/journal.pcbi.1002822>.
- Cantor, R.M., Lange, K. & Sinsheimer, J.S., 2010. Prioritizing GWAS results: A review of statistical methods and recommendations for their application. *American journal of human genetics*, 86(1), pp.6–22. Available at: <http://dx.doi.org/10.1016/j.ajhg.2009.11.017>.
- Chen, Y. et al., 2012. Systematic evaluation of factors influencing ChIP-seq fidelity. *Nature methods*, 9(6), pp.609–614. Available at: <http://dx.doi.org/10.1038/nmeth.1985>.
- Cock, P.J.A. et al., 2010. The Sanger FASTQ file format for sequences with quality scores, and the Solexa/Illumina FASTQ variants. *Nucleic acids research*, 38(6), pp.1767–1771. Available at: <http://dx.doi.org/10.1093/nar/gkp1137>.
- Colombel, J.-F. et al., 2007. Adalimumab for maintenance of clinical response and remission in patients with Crohn's disease: the CHARM trial. *Gastroenterology*, 132(1), pp.52–65. Available at: <http://dx.doi.org/10.1053/j.gastro.2006.11.041>.
- Dale, R.K., Pedersen, B.S. & Quinlan, A.R., 2011. Pybedtools: a flexible Python library for manipulating genomic datasets and annotations. *Bioinformatics*, 27(24), pp.3423–3424. Available at:

<http://dx.doi.org/10.1093/bioinformatics/btr539>.

- Das, R. et al., 2006. Computational prediction of methylation status in human genomic sequences. *Proceedings of the National Academy of Sciences of the United States of America*, 103(28), pp.10713–10716. Available at: <http://dx.doi.org/10.1073/pnas.0602949103>.
- Eddy, S.R., 1998. Profile hidden Markov models. *Bioinformatics*, 14(9), pp.755–763. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9918945>.
- Edwards, S.L. et al., 2013. Beyond GWASs: illuminating the dark road from association to function. *American journal of human genetics*, 93(5), pp.779–797. Available at: <http://dx.doi.org/10.1016/j.ajhg.2013.10.012>.
- Encinas, J.A. et al., 1999. QTL influencing autoimmune diabetes and encephalomyelitis map to a 0.15-cM region containing Il2. *Nature genetics*, 21(2), pp.158–160. Available at: <http://dx.doi.org/10.1038/5941>.
- ENCODE Project Consortium, 2012. An integrated encyclopedia of DNA elements in the human genome. *Nature*, 489(7414), pp.57–74. Available at: <http://dx.doi.org/10.1038/nature11247>.
- ENCODE Project Consortium et al., 2007. Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project. *Nature*, 447(7146), pp.799–816. Available at: <http://dx.doi.org/10.1038/nature05874>.
- ENCODE Project Consortium, 2004. The ENCODE (ENCyclopedia Of DNA Elements) Project. *Science*, 306(5696), pp.636–640. Available at: <http://dx.doi.org/10.1126/science.1105136>.
- Ernst, J. & Kellis, M., 2010. Discovery and characterization of chromatin states for systematic annotation of the human genome. *Nature biotechnology*, 28(8), pp.817–825. Available at: <http://dx.doi.org/10.1038/nbt.1662>.
- Ernst, J. & Kellis, M., 2015. Large-scale imputation of epigenomic datasets for systematic annotation of diverse human tissues. *Nature biotechnology*, 33, p.364. Available at: <http://dx.doi.org/10.1038/nbt.3157>.
- Evans, J.P. et al., 2011. Genomics. Deflating the genomic bubble. *Science*, 331(6019), pp.861–862. Available at: <http://dx.doi.org/10.1126/science.1198039>.
- Fletcher, J.M. et al., 2009. CD39+Foxp3+ regulatory T Cells suppress pathogenic Th17 cells and are impaired in multiple sclerosis. *Journal of immunology*, 183(11), pp.7602–7610. Available at: <http://dx.doi.org/10.4049/jimmunol.0901881>.
- Freedman, M.L. et al., 2011. Principles for the post-GWAS functional characterization of cancer risk loci. *Nature genetics*, 43(6), pp.513–518.

Available at: <http://dx.doi.org/10.1038/ng.840>.

- Furst, D.E., 2010. The risk of infections with biologic therapies for rheumatoid arthritis. *Seminars in arthritis and rheumatism*, 39(5), pp.327–346. Available at: <http://dx.doi.org/10.1016/j.semarthrit.2008.10.002>.
- Gabriel, S.B. et al., 2002. The structure of haplotype blocks in the human genome. *Science*, 296(5576), pp.2225–2229. Available at: <http://dx.doi.org/10.1126/science.1069424>.
- Gates, L.A. et al., 2017. Acetylation on histone H3 lysine 9 mediates a switch from transcription initiation to elongation. *The Journal of biological chemistry*. Available at: <http://www.jbc.org/content/early/2017/07/17/jbc.M117.802074.abstract>.
- Guerini, F.R. et al., 2012. A functional variant in ERAP1 predisposes to multiple sclerosis. *PloS one*, 7(1), p.e29931. Available at: <http://dx.doi.org/10.1371/journal.pone.0029931>.
- Hoffmann, T.J. & Witte, J.S., 2015. Strategies for Imputing and Analyzing Rare Variants in Association Studies. *Trends in genetics: TIG*, 31(10), pp.556–563. Available at: <http://dx.doi.org/10.1016/j.tig.2015.07.006>.
- Holland, S.M. et al., 2007. STAT3 mutations in the hyper-IgE syndrome. *The New England journal of medicine*, 357(16), pp.1608–1619. Available at: <http://dx.doi.org/10.1056/NEJMoa073687>.
- Howie, B. et al., 2012. Fast and accurate genotype imputation in genome-wide association studies through pre-phasing. *Nature genetics*, 44(8), pp.955–959. Available at: <http://dx.doi.org/10.1038/ng.2354>.
- Howie, B.N., Donnelly, P. & Marchini, J., 2009. A flexible and accurate genotype imputation method for the next generation of genome-wide association studies. *PLoS genetics*, 5(6), p.e1000529. Available at: <http://dx.doi.org/10.1371/journal.pgen.1000529>.
- Hrdlickova, B. et al., 2014. Genetic variation in the non-coding genome: Involvement of micro-RNAs and long non-coding RNAs in disease. *Biochimica et biophysica acta*, 1842(10), pp.1910–1922. Available at: <http://dx.doi.org/10.1016/j.bbadis.2014.03.011>.
- Hsu, F. et al., 2006. The UCSC Known Genes. *Bioinformatics*, 22(9), pp.1036–1046. Available at: <https://academic.oup.com/bioinformatics/article/22/9/1036/200093> [Accessed August 1, 2018].
- Hunter, D.J. & Kraft, P., 2007. Drinking from the fire hose--statistical issues in genomewide association studies. *The New England journal of medicine*, 357(5),

- pp.436–439. Available at: <http://dx.doi.org/10.1056/NEJMp078120>.
- Hunter, J.D., 2007. Matplotlib: A 2D Graphics Environment. *Computing in Science Engineering*, 9(3), pp.90–95. Available at: <http://dx.doi.org/10.1109/MCSE.2007.55>.
- Jerez, J.M. et al., 2010. Missing data imputation using statistical and machine learning methods in a real breast cancer problem. *Artificial intelligence in medicine*, 50(2), pp.105–115. Available at: <http://dx.doi.org/10.1016/j.artmed.2010.05.002>.
- Jung, Y.L. et al., 2014. Impact of sequencing depth in ChIP-seq experiments. *Nucleic acids research*, 42(9), p.e74. Available at: <http://dx.doi.org/10.1093/nar/gku178>.
- Karlič, R. et al., 2010. Histone modification levels are predictive for gene expression. *Proceedings of the National Academy of Sciences of the United States of America*, 107(7), pp.2926–2931. Available at: <http://www.pnas.org/content/107/7/2926> [Accessed August 21, 2018].
- Kent, W.J. et al., 2002. The human genome browser at UCSC. *Genome research*, 12(6), pp.996–1006. Available at: <http://dx.doi.org/10.1101/gr.229102>.
- Kluyver, T. et al., 2016. Jupyter Notebooks – a publishing format for reproducible computational workflows. In F. Loizides & B. Schmidt, eds. *Positioning and Power in Academic Publishing: Players, Agents and Agendas*. IOS Press, pp. 87–90. Available at: <https://eprints.soton.ac.uk/403913/> [Accessed August 1, 2018].
- Korber, P. et al., 2004. Evidence for histone eviction in trans upon induction of the yeast PHO5 promoter. *Molecular and cellular biology*, 24(24), pp.10965–10974. Available at: <http://dx.doi.org/10.1128/MCB.24.24.10965-10974.2004>.
- Korte, A. & Farlow, A., 2013. The advantages and limitations of trait analysis with GWAS: a review. *Plant methods*, 9, p.29. Available at: <http://dx.doi.org/10.1186/1746-4811-9-29>.
- Kouzarides, T., 2007. Chromatin modifications and their function. *Cell*, 128(4), pp.693–705. Available at: <http://dx.doi.org/10.1016/j.cell.2007.02.005>.
- Kriegler, M. et al., 1988. A novel form of TNF/cachectin is a cell surface cytotoxic transmembrane protein: ramifications for the complex physiology of TNF. *Cell*, 53(1), pp.45–53. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/3349526>.
- Krogh, A., 1998. An Introduction to Hidden Markov Models for Biological Sequences. In D. B. S. A. S. K. S. L. Salzberg, ed. *Computational Methods in Molecular Biology*. Elsevier, pp. 45–63. Available at: <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.85.7972&rep=rep1&ty>

pe=pdf.

- Kronenberg, M. & Rudensky, A., 2005. Regulation of immunity by self-reactive T cells. *Nature*, 435(7042), pp.598–604. Available at: <http://dx.doi.org/10.1038/nature03725>.
- Landt, S.G. et al., 2012. ChIP-seq guidelines and practices of the ENCODE and modENCODE consortia. *Genome research*, 22(9), pp.1813–1831. Available at: <http://dx.doi.org/10.1101/gr.136184.111>.
- Lauberth, S.M. et al., 2013. H3K4me3 interactions with TAF3 regulate preinitiation complex assembly and selective gene activation. *Cell*, 152(5), pp.1021–1036. Available at: <http://dx.doi.org/10.1016/j.cell.2013.01.052>.
- Liang, G. et al., 2004. Distinct localization of histone H3 acetylation and H3-K4 methylation to the transcription start sites in the human genome. *Proceedings of the National Academy of Sciences of the United States of America*, 101(19), pp.7357–7362. Available at: <http://dx.doi.org/10.1073/pnas.0401866101>.
- Librado, P. & Rozas, J., 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics*, 25(11), pp.1451–1452. Available at: <http://dx.doi.org/10.1093/bioinformatics/btp187>.
- Li, H., 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv [q-bio.GN]*. Available at: <http://arxiv.org/abs/1303.3997>.
- Li, H. et al., 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, 25(16), pp.2078–2079. Available at: <http://dx.doi.org/10.1093/bioinformatics/btp352>.
- Liu, J. et al., 2015. Chromatin Landscape Defined by Repressive Histone Methylation during Oligodendrocyte Differentiation. *The Journal of neuroscience: the official journal of the Society for Neuroscience*, 35(1), pp.352–365. Available at: <http://www.jneurosci.org/content/35/1/352> [Accessed August 23, 2018].
- Li, X. & Zheng, Y., 2015. Regulatory T cell identity: formation and maintenance. *Trends in immunology*, 36(6), pp.344–353. Available at: <http://dx.doi.org/10.1016/j.it.2015.04.006>.
- Li, Y. et al., 2009. Genotype imputation. *Annual review of genomics and human genetics*, 10, pp.387–406. Available at: <http://dx.doi.org/10.1146/annurev.genom.9.081307.164242>.
- Lorch, Y., Maier-Davis, B. & Kornberg, R.D., 2010. Mechanism of chromatin remodeling. *Proceedings of the National Academy of Sciences of the United States of America*, 107(8), pp.3458–3462. Available at:

<http://dx.doi.org/10.1073/pnas.1000398107>.

- Manolio, T.A., 2013. Bringing genome-wide association findings into clinical use. *Nature reviews. Genetics*, 14, p.549. Available at: <http://dx.doi.org/10.1038/nrg3523>.
- Manolio, T.A. et al., 2009. Finding the missing heritability of complex diseases. *Nature*, 461(7265), pp.747–753. Available at: <http://dx.doi.org/10.1038/nature08494>.
- Marchini, J. & Howie, B., 2010. Genotype imputation for genome-wide association studies. *Nature reviews. Genetics*, 11(7), pp.499–511. Available at: <http://dx.doi.org/10.1038/nrg2796>.
- Martin, E.R. et al., 2000. SNPing away at complex diseases: analysis of single-nucleotide polymorphisms around APOE in Alzheimer disease. *American journal of human genetics*, 67(2), pp.383–394. Available at: <http://dx.doi.org/10.1086/303003>.
- McCarthy, S. et al., 2016. A reference panel of 64,976 haplotypes for genotype imputation. *Nature genetics*, 48(10), pp.1279–1283. Available at: <http://dx.doi.org/10.1038/ng.3643>.
- McKerns, M.M. et al., 2012. Building a Framework for Predictive Science. *arXiv [cs.MS]*. Available at: <http://arxiv.org/abs/1202.1056>.
- Mc Kinney, W., pandas: a Foundational Python Library for Data Analysis and Statistics. Available at: https://www.dlr.de/sc/Portaldata/15/Resources/dokumente/pyhpc2011/submissions/pyhpc2011_submission_9.pdf.
- McLean, C.Y. et al., 2010. GREAT improves functional interpretation of cis-regulatory regions. *Nature biotechnology*, 28(5), pp.495–501. Available at: <http://dx.doi.org/10.1038/nbt.1630>.
- Mertes, F. et al., 2011. Targeted enrichment of genomic DNA regions for next-generation sequencing. *Briefings in functional genomics*, 10(6), pp.374–386. Available at: <https://academic.oup.com/bfg/article/10/6/374/233492> [Accessed August 20, 2018].
- Mikkelsen, T.S. et al., 2007. Genome-wide maps of chromatin state in pluripotent and lineage-committed cells. *Nature*, 448(7153), pp.553–560. Available at: <http://dx.doi.org/10.1038/nature06008>.
- Millman, K.J. & Aivazis, M., 2011. Python for Scientists and Engineers. *Computing in Science Engineering*, 13(2), pp.9–12. Available at: <http://dx.doi.org/10.1109/MCSE.2011.36>.

- NCI-NHGRI Working Group on Replication in Association Studies et al., 2007. Replicating genotype-phenotype associations. *Nature*, 447(7145), pp.655–660. Available at: <http://dx.doi.org/10.1038/447655a>.
- O'Geen, H., Echipare, L. & Farnham, P.J., 2011. Using ChIP-seq technology to generate high-resolution profiles of histone modifications. *Methods in molecular biology*, 791, pp.265–286. Available at: http://dx.doi.org/10.1007/978-1-61779-316-5_20.
- Oliphant, T.E., 2006. *A guide to NumPy*, Trelgol Publishing USA. Available at: <http://web.mit.edu/dvp/Public/numpybook.pdf>.
- Oliphant, T.E., 2007. Python for Scientific Computing. *Computing in Science Engineering*, 9(3), pp.10–20. Available at: <http://dx.doi.org/10.1109/MCSE.2007.58>.
- Park, P.J., 2009. ChIP-seq: advantages and challenges of a maturing technology. *Nature reviews. Genetics*, 10(10), pp.669–680. Available at: <http://dx.doi.org/10.1038/nrg2641>.
- Pedregosa, F. et al., 2011. Scikit-learn: Machine Learning in Python. *Journal of machine learning research: JMLR*, 12(Oct), pp.2825–2830. Available at: <http://www.jmlr.org/papers/volume12/pedregosa11a/pedregosa11a.pdf> [Accessed August 1, 2018].
- Pei, Y.-F. et al., 2010. Analyses and comparison of imputation-based association methods. *PloS one*, 5(5), p.e10827. Available at: <http://dx.doi.org/10.1371/journal.pone.0010827>.
- Perez, F. & Granger, B.E., 2007. IPython: A System for Interactive Scientific Computing. *Computing in Science Engineering*, 9(3), pp.21–29. Available at: <http://dx.doi.org/10.1109/MCSE.2007.53>.
- Pidasheva, S. et al., 2011. Functional studies on the IBD susceptibility gene IL23R implicate reduced receptor function in the protective genetic variant R381Q. *PloS one*, 6(10), p.e25038. Available at: <http://dx.doi.org/10.1371/journal.pone.0025038>.
- Pierini, A. et al., 2014. CD4+FoxP3+ Regulatory T Cells Regulate B Cell Differentiation and Induce Tolerance to Bone Marrow Grafts. *Blood*, 124(21), pp.4322–4322. Available at: <http://www.bloodjournal.org/content/124/21/4322?sso-checked=true> [Accessed August 19, 2018].
- Portales-Casamar, E. et al., 2010. JASPAR 2010: the greatly expanded open-access database of transcription factor binding profiles. *Nucleic acids research*, 38(Database issue), pp.D105–10. Available at:

<http://dx.doi.org/10.1093/nar/gkp950>.

Purcell, S. et al., 2007. PLINK: a tool set for whole-genome association and population-based linkage analyses. *American journal of human genetics*, 81(3), pp.559–575. Available at: <http://dx.doi.org/10.1086/519795>.

Quail, M.A. et al., 2008. A large genome center's improvements to the Illumina sequencing system. *Nature methods*, 5(12), pp.1005–1010. Available at: <http://dx.doi.org/10.1038/nmeth.1270>.

Quinlan, A.R. & Hall, I.M., 2010. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics*, 26(6), pp.841–842. Available at: <http://dx.doi.org/10.1093/bioinformatics/btq033>.

Radman-Livaja, M. & Rando, O.J., 2010. Nucleosome positioning: how is it established, and why does it matter? *Developmental biology*, 339(2), pp.258–266. Available at: <http://dx.doi.org/10.1016/j.ydbio.2009.06.012>.

Ramos, R.G. & Olden, K., 2008. Gene-environment interactions in the development of complex disease phenotypes. *International journal of environmental research and public health*, 5(1), pp.4–11. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18441400>.

Raychaudhuri, S. et al., 2012. Five amino acids in three HLA proteins explain most of the association between MHC and seropositive rheumatoid arthritis. *Nature genetics*, 44(3), pp.291–296. Available at: <http://dx.doi.org/10.1038/ng.1076>.

Raychaudhuri, S., 2011. Mapping rare and common causal alleles for complex human diseases. *Cell*, 147(1), pp.57–69. Available at: <http://dx.doi.org/10.1016/j.cell.2011.09.011>.

Rea, S. et al., 2000. Regulation of chromatin structure by site-specific histone H3 methyltransferases. *Nature*, 406(6796), pp.593–599. Available at: <http://dx.doi.org/10.1038/35020506>.

Rioux, J.D. & Abbas, A.K., 2005. Paths to understanding the genetic basis of autoimmune disease. *Nature*, 435(7042), pp.584–589. Available at: <http://dx.doi.org/10.1038/nature03723>.

Rivera, C.M. & Ren, B., 2013. Mapping Human Epigenomes. *Cell*, 155(1), pp.39–55. Available at: [https://www.cell.com/fulltext/S0092-8674\(13\)01148-3](https://www.cell.com/fulltext/S0092-8674(13)01148-3) [Accessed August 21, 2018].

Robertson, G. et al., 2007. Genome-wide profiles of STAT1 DNA association using chromatin immunoprecipitation and massively parallel sequencing. *Nature methods*, 4(8), pp.651–657. Available at: <http://dx.doi.org/10.1038/nmeth1068>.

Rosenblum, M.D., Remedios, K.A. & Abbas, A.K., 2015. Mechanisms of human

- autoimmunity. *The Journal of clinical investigation*, 125(6), pp.2228–2233. Available at: <http://dx.doi.org/10.1172/JCI78088>.
- Rosenfeld, J.A. et al., 2009. Determination of enriched histone modifications in non-genic portions of the human genome. *BMC genomics*, 10, p.143. Available at: <http://dx.doi.org/10.1186/1471-2164-10-143>.
- Roshyara, N.R. et al., 2016. Comparing performance of modern genotype imputation methods in different ethnicities. *Scientific reports*, 6, p.34386. Available at: <http://dx.doi.org/10.1038/srep34386>.
- Rutgeerts, P. et al., 2005. Infliximab for induction and maintenance therapy for ulcerative colitis. *The New England journal of medicine*, 353(23), pp.2462–2476. Available at: <http://dx.doi.org/10.1056/NEJMoa050516>.
- Sakaguchi, S. et al., 1995. Immunologic self-tolerance maintained by activated T cells expressing IL-2 receptor alpha-chains (CD25). Breakdown of a single mechanism of self-tolerance causes various autoimmune diseases. *Journal of immunology*, 155(3), pp.1151–1164. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/7636184>.
- Sakaguchi, S. et al., 2008. Regulatory T cells and immune tolerance. *Cell*, 133(5), pp.775–787. Available at: <http://dx.doi.org/10.1016/j.cell.2008.05.009>.
- Schmidl, C. et al., 2015. ChIPmentation: fast, robust, low-input ChIP-seq for histones and transcription factors. *Nature methods*, 12(10), pp.963–965. Available at: <http://dx.doi.org/10.1038/nmeth.3542>.
- Schones, D.E. et al., 2008. Dynamic regulation of nucleosome positioning in the human genome. *Cell*, 132(5), pp.887–898. Available at: <http://dx.doi.org/10.1016/j.cell.2008.02.022>.
- Schork, A.J. et al., 2013. All SNPs are not created equal: genome-wide association studies reveal a consistent pattern of enrichment among functionally annotated SNPs. *PLoS genetics*, 9(4), p.e1003449. Available at: <http://dx.doi.org/10.1371/journal.pgen.1003449>.
- Shu, W. et al., 2011. Genome-wide analysis of the relationships between DNaseI HS, histone modifications and gene expression reveals distinct modes of chromatin domains. *Nucleic acids research*, 39(17), pp.7428–7443. Available at: <http://dx.doi.org/10.1093/nar/gkr443>.
- Spain, S.L. & Barrett, J.C., 2015. Strategies for fine-mapping complex traits. *Human molecular genetics*, 24(R1), pp.R111–9. Available at: <http://dx.doi.org/10.1093/hmg/ddv260>.
- Spencer, C.C.A. et al., 2009. Designing genome-wide association studies: sample size, power, imputation, and the choice of genotyping chip. *PLoS genetics*, 5(5),

p.e1000477. Available at: <http://dx.doi.org/10.1371/journal.pgen.1000477>.

Starmer, J. & Magnuson, T., 2016. Detecting broad domains and narrow peaks in ChIP-seq data with hiddenDomains. *BMC bioinformatics*, 17, p.144. Available at: <http://dx.doi.org/10.1186/s12859-016-0991-z>.

Stunnenberg, H.G. et al., 2016. The International Human Epigenome Consortium: A Blueprint for Scientific Collaboration and Discovery. *Cell*, 167(5), pp.1145–1149. Available at: [https://www.cell.com/cell/abstract/S0092-8674\(16\)31528-8](https://www.cell.com/cell/abstract/S0092-8674(16)31528-8) [Accessed August 18, 2018].

Tehranchi, A.K. et al., 2016. Pooled ChIP-Seq Links Variation in Transcription Factor Binding to Complex Disease Risk. *Cell*, 165(3), pp.730–741. Available at: <http://dx.doi.org/10.1016/j.cell.2016.03.041>.

The International HapMap 3 Consortium et al., 2010. Integrating common and rare genetic variation in diverse human populations. *Nature*, 467, p.52. Available at: <http://dx.doi.org/10.1038/nature09298>.

Thurman, R.E. et al., 2012. The accessible chromatin landscape of the human genome. *Nature*, 489(7414), pp.75–82. Available at: <http://dx.doi.org/10.1038/nature11232>.

Troyanskaya, O. et al., 2001. Missing value estimation methods for DNA microarrays. *Bioinformatics*, 17(6), pp.520–525. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/11395428>.

Trynka, G. et al., 2013. Chromatin marks identify critical cell types for fine mapping complex trait variants. *Nature genetics*, 45(2), pp.124–130. Available at: <http://dx.doi.org/10.1038/ng.2504>.

Trynka, G. & Raychaudhuri, S., 2013. Using chromatin marks to interpret and localize genetic associations to complex human traits and diseases. *Current opinion in genetics & development*, 23(6), pp.635–641. Available at: <http://dx.doi.org/10.1016/j.gde.2013.10.009>.

Tsompana, M. & Buck, M.J., 2014. Chromatin accessibility: a window into the genome. *Epigenetics & chromatin*, 7(1), p.33. Available at: <https://doi.org/10.1186/1756-8935-7-33>.

Valouev, A. et al., 2008. Genome-wide analysis of transcription factor binding sites based on ChIP-Seq data. *Nature methods*, 5(9), pp.829–834. Available at: <http://dx.doi.org/10.1038/nmeth.1246>.

Varmus, H., 2010. Ten years on--the human genome and medicine. *The New England journal of medicine*, 362(21), pp.2028–2029. Available at: <http://dx.doi.org/10.1056/NEJMe0911933>.

- Visel, A. et al., 2009. ChIP-seq accurately predicts tissue-specific activity of enhancers. *Nature*, 457(7231), pp.854–858. Available at: <http://dx.doi.org/10.1038/nature07730>.
- Visscher, P.M. et al., 2017. 10 Years of GWAS Discovery: Biology, Function, and Translation. *American journal of human genetics*, 101(1), pp.5–22. Available at: <http://dx.doi.org/10.1016/j.ajhg.2017.06.005>.
- Wang, L. et al., 2014. Breach of tolerance: primary biliary cirrhosis. *Seminars in liver disease*, 34(3), pp.297–317. Available at: <http://dx.doi.org/10.1055/s-0034-1383729>.
- Waskom, M. et al., 2014. *seaborn: v0.5.0 (November 2014)*, Available at: <https://zenodo.org/record/12710>.
- Wellcome Trust Case Control Consortium et al., 2010. Genome-wide association study of CNVs in 16,000 cases of eight common diseases and 3,000 shared controls. *Nature*, 464(7289), pp.713–720. Available at: <http://dx.doi.org/10.1038/nature08979>.
- Yu, G., Wang, L.-G. & He, Q.-Y., 2015. ChIPseeker: an R/Bioconductor package for ChIP peak annotation, comparison and visualization. *Bioinformatics*, 31(14), pp.2382–2383. Available at: <http://dx.doi.org/10.1093/bioinformatics/btv145>.
- Zhang, F. & Lupski, J.R., 2015. Non-coding genetic variants in human disease. *Human molecular genetics*, 24(R1), pp.R102–10. Available at: <http://dx.doi.org/10.1093/hmg/ddv259>.
- Zhang, Y. et al., 2008. Model-based Analysis of ChIP-Seq (MACS). *Genome biology*, 9(9), p.R137. Available at: <https://doi.org/10.1186/gb-2008-9-9-r137>.
- Zhang, Z. & Pugh, B.F., 2011. High-resolution genome-wide mapping of the primary structure of chromatin. *Cell*, 144(2), pp.175–186. Available at: <http://dx.doi.org/10.1016/j.cell.2011.01.003>.
- Zhou, J. et al., 2011. A20-binding inhibitor of NF- κ B (ABIN1) controls Toll-like receptor-mediated CCAAT/enhancer-binding protein β activation and protects from inflammatory disease. *Proceedings of the National Academy of Sciences of the United States of America*, 108(44), pp.E998–1006. Available at: <http://dx.doi.org/10.1073/pnas.1106232108>.
- Zhou, J. & Troyanskaya, O.G., 2015. Predicting effects of noncoding variants with deep learning-based sequence model. *Nature methods*, 12(10), pp.931–934. Available at: <http://dx.doi.org/10.1038/nmeth.3547>.