

REFERENCES

- Abell, K., Bilancio, A., Clarkson, R.W., Tiffen, P.G., Altaparmakov, A.I., Burdon, T.G., Asano, T., Vanhaesebroeck, B., and Watson, C.J. (2005). Stat3-induced apoptosis requires a molecular switch in PI(3)K subunit composition. *Nat Cell Biol* 7, 392-398.
- Alvi, A.J., Clayton, H., Joshi, C., Enver, T., Ashworth, A., Vivanco, M.M., Dale, T.C., and Smalley, M.J. (2003). Functional and molecular characterisation of mammary side population cells. *Breast Cancer Res* 5, R1-8.
- Amsen, D., Antov, A., Jankovic, D., Sher, A., Radtke, F., Souabni, A., Busslinger, M., McCright, B., Gridley, T., and Flavell, R.A. (2007). Direct regulation of Gata3 expression determines the T helper differentiation potential of Notch. *Immunity* 27, 89-99.
- Angrand, P.O., Daigle, N., van der Hoeven, F., Scholer, H.R., and Stewart, A.F. (1999). Simplified generation of targeting constructs using ET recombination. *Nucleic Acids Res* 27, e16.
- Ansel, K.M., Djuretic, I., Tanasa, B., and Rao, A. (2006). Regulation of Th2 differentiation and Il4 locus accessibility. *Annual review of immunology* 24, 607-656.
- Artavanis-Tsakonas, S., Rand, M.D., and Lake, R.J. (1999). Notch signaling: cell fate control and signal integration in development. *Science (New York, NY)* 284, 770-776.
- Asselin-Labat, M.L., Shackleton, M., Stingl, J., Vaillant, F., Forrest, N.C., Eaves, C.J., Visvader, J.E., and Lindeman, G.J. (2006). Steroid hormone receptor status of mouse mammary stem cells. *Journal of the National Cancer Institute* 98, 1011-1014.
- Asselin-Labat, M.L., Sutherland, K.D., Barker, H., Thomas, R., Shackleton, M., Forrest, N.C., Hartley, L., Robb, L., Grosveld, F.G., van der Wees, J., *et al.* (2007). Gata-3 is an essential regulator of mammary-gland morphogenesis and luminal-cell differentiation. *Nat Cell Biol* 9, 201-209.
- Austin, C.P., Battey, J.F., Bradley, A., Bucan, M., Capecchi, M., Collins, F.S., Dove, W.F., Duyk, G., Dymecki, S., Eppig, J.T., *et al.* (2004). The knockout mouse project. *Nature genetics* 36, 921-924.
- Auwerx, J., Avner, P., Baldock, R., Ballabio, A., Balling, R., Barbacid, M., Berns, A., Bradley, A., Brown, S., Carmeliet, P., *et al.* (2004). The European dimension for the mouse genome mutagenesis program. *Nature genetics* 36, 925-927.
- Avram, D., Fields, A., Pretty On Top, K., Nevriy, D.J., Ishmael, J.E., and Leid, M. (2000). Isolation of a novel family of C(2)H(2) zinc finger proteins implicated in transcriptional repression mediated by chicken ovalbumin upstream promoter transcription factor (COUP-TF) orphan nuclear receptors. *J Biol Chem* 275, 10315-10322.
- Avram, D., Fields, A., Senawong, T., Topark-Ngarm, A., and Leid, M. (2002). COUP-TF (chicken ovalbumin upstream promoter transcription factor)-interacting protein 1 (CTIP1) is a sequence-specific DNA binding protein. *Biochem J* 368, 555-563.
- Behrens, J., von Kries, J.P., Kuhl, M., Bruhn, L., Wedlich, D., Grosschedl, R., and Birchmeier, W. (1996). Functional interaction of beta-catenin with the transcription factor LEF-1. *Nature* 382, 638-642.
- Bocchinfuso, W.P., and Korach, K.S. (1997). Mammary gland development and tumorigenesis in estrogen receptor knockout mice. *J Mammary Gland Biol Neoplasia* 2, 323-334.

- Booth, B.W., and Smith, G.H. (2006). Estrogen receptor-alpha and progesterone receptor are expressed in label-retaining mammary epithelial cells that divide asymmetrically and retain their template DNA strands. *Breast Cancer Res* 8, R49.
- Boras-Granic, K., Chang, H., Grosschedl, R., and Hamel, P.A. (2006). Lef1 is required for the transition of Wnt signaling from mesenchymal to epithelial cells in the mouse embryonic mammary gland. *Developmental biology* 295, 219-231.
- Bradley, A., Evans, M., Kaufman, M.H., and Robertson, E. (1984). Formation of germ-line chimaeras from embryo-derived teratocarcinoma cell lines. *Nature* 309, 255-256.
- Bradley, A., and Liu, P. (1996). Target practice in transgenics. *Nature genetics* 14, 121-123.
- Brinster, R.L. (1974). The effect of cells transferred into the mouse blastocyst on subsequent development. *The Journal of experimental medicine* 140, 1049-1056.
- Brinster, R.L., Chen, H.Y., Trumbauer, M., Senechal, A.W., Warren, R., and Palmiter, R.D. (1981). Somatic expression of herpes thymidine kinase in mice following injection of a fusion gene into eggs. *Cell* 27, 223-231.
- Brisken, C., Heineman, A., Chavarria, T., Elenbaas, B., Tan, J., Dey, S.K., McMahon, J.A., McMahon, A.P., and Weinberg, R.A. (2000). Essential function of Wnt-4 in mammary gland development downstream of progesterone signaling. *Genes Dev* 14, 650-654.
- Bryder, D., Rossi, D.J., and Weissman, I.L. (2006). Hematopoietic stem cells: the paradigmatic tissue-specific stem cell. *The American journal of pathology* 169, 338-346.
- Buchholz, F., Angrand, P.O., and Stewart, A.F. (1998). Improved properties of FLP recombinase evolved by cycling mutagenesis. *Nature biotechnology* 16, 657-662.
- Buono, K.D., Robinson, G.W., Martin, C., Shi, S., Stanley, P., Tanigaki, K., Honjo, T., and Hennighausen, L. (2006). The canonical Notch/RBP-J signaling pathway controls the balance of cell lineages in mammary epithelium during pregnancy. *Developmental biology* 293, 565-580.
- Cantor, A.B., and Orkin, S.H. (2001). Hematopoietic development: a balancing act. *Current opinion in genetics & development* 11, 513-519.
- Capecchi, M.R. (1989). Altering the genome by homologous recombination. *Science (New York, NY)* 244, 1288-1292.
- Carroll, J.S., Liu, X.S., Brodsky, A.S., Li, W., Meyer, C.A., Szary, A.J., Eeckhoutte, J., Shao, W., Hestermann, E.V., Geistlinger, T.R., *et al.* (2005). Chromosome-wide mapping of estrogen receptor binding reveals long-range regulation requiring the forkhead protein FoxA1. *Cell* 122, 33-43.
- Chan, W., Costantino, N., Li, R., Lee, S.C., Su, Q., Melvin, D., Court, D.L., and Liu, P. (2007). A recombineering based approach for high-throughput conditional knockout targeting vector construction. *Nucleic Acids Res* 35, e64.
- Chapman, R.S., Lourenco, P., Tonner, E., Flint, D., Selbert, S., Takeda, K., Akira, S., Clarke, A.R., and Watson, C.J. (2000). The role of Stat3 in apoptosis and mammary gland involution. Conditional deletion of Stat3. *Adv Exp Med Biol* 480, 129-138.
- Chapman, R.S., Lourenco, P.C., Tonner, E., Flint, D.J., Selbert, S., Takeda, K., Akira, S., Clarke, A.R., and Watson, C.J. (1999). Suppression of epithelial apoptosis and delayed mammary gland involution in mice with a conditional knockout of Stat3. *Genes Dev* 13, 2604-2616.

- Chu, E.Y., Hens, J., Andl, T., Kairo, A., Yamaguchi, T.P., Brisken, C., Glick, A., Wysolmerski, J.J., and Millar, S.E. (2004). Canonical WNT signaling promotes mammary placode development and is essential for initiation of mammary gland morphogenesis. *Development (Cambridge, England)* *131*, 4819-4829.
- Cismasiu, V.B., Adamo, K., Gecewicz, J., Duque, J., Lin, Q., and Avram, D. (2005). BCL11B functionally associates with the NuRD complex in T lymphocytes to repress targeted promoter. *Oncogene* *24*, 6753-6764.
- Clarke, R.B., Howell, A., Potten, C.S., and Anderson, E. (1997). Dissociation between steroid receptor expression and cell proliferation in the human breast. *Cancer research* *57*, 4987-4991.
- Collins, F.S., Rossant, J., and Wurst, W. (2007). A mouse for all reasons. *Cell* *128*, 9-13.
- Copeland, N.G., Jenkins, N.A., and Court, D.L. (2001). Recombineering: a powerful new tool for mouse functional genomics. *Nat Rev Genet* *2*, 769-779.
- Costa, R.H., Kalinichenko, V.V., and Lim, L. (2001). Transcription factors in mouse lung development and function. *American journal of physiology* *280*, L823-838.
- Covarrubias, L., Cervantes, L., Covarrubias, A., Soberon, X., Vichido, I., Blanco, A., Kupersztoch-Portnoy, Y.M., and Bolivar, F. (1981). Construction and characterization of new cloning vehicles. V. Mobilization and coding properties of pBR322 and several deletion derivatives including pBR327 and pBR328. *Gene* *13*, 25-35.
- Cui, Y., Riedlinger, G., Miyoshi, K., Tang, W., Li, C., Deng, C.X., Robinson, G.W., and Hennighausen, L. (2004). Inactivation of Stat5 in mouse mammary epithelium during pregnancy reveals distinct functions in cell proliferation, survival, and differentiation. *Mol Cell Biol* *24*, 8037-8047.
- Cunha, G.R., and Hom, Y.K. (1996). Role of mesenchymal-epithelial interactions in mammary gland development. *J Mammary Gland Biol Neoplasia* *1*, 21-35.
- Daniel, C.W., Silberstein, G.B., and Strickland, P. (1987). Direct action of 17 beta-estradiol on mouse mammary ducts analyzed by sustained release implants and steroid autoradiography. *Cancer research* *47*, 6052-6057.
- Darnell, J.E., Jr. (1997). STATs and gene regulation. *Science (New York, NY)* *277*, 1630-1635.
- Datsenko, K.A., and Wanner, B.L. (2000). One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proceedings of the National Academy of Sciences of the United States of America* *97*, 6640-6645.
- Datta, S., Costantino, N., and Court, D.L. (2006). A set of recombineering plasmids for gram-negative bacteria. *Gene* *379*, 109-115.
- Davis, M.M. (2007). Blimp-1 over Budapest. *Nat Immunol* *8*, 445-447.
- Deng, C., and Capecchi, M.R. (1992). Reexamination of gene targeting frequency as a function of the extent of homology between the targeting vector and the target locus. *Mol Cell Biol* *12*, 3365-3371.
- Deome, K.B., Faulkin, L.J., Jr., Bern, H.A., and Blair, P.B. (1959). Development of mammary tumors from hyperplastic alveolar nodules transplanted into gland-free mammary fat pads of female C3H mice. *Cancer research* *19*, 515-520.
- Donehower, L.A., Harvey, M., Slagle, B.L., McArthur, M.J., Montgomery, C.A., Jr., Butel, J.S., and Bradley, A. (1992). Mice deficient for p53 are developmentally normal but susceptible to spontaneous tumours. *Nature* *356*, 215-221.

- Dontu, G., Abdallah, W.M., Foley, J.M., Jackson, K.W., Clarke, M.F., Kawamura, M.J., and Wicha, M.S. (2003). In vitro propagation and transcriptional profiling of human mammary stem/progenitor cells. *Genes Dev* 17, 1253-1270.
- Dontu, G., El-Ashry, D., and Wicha, M.S. (2004). Breast cancer, stem/progenitor cells and the estrogen receptor. *Trends in endocrinology and metabolism: TEM* 15, 193-197.
- Eblaghie, M.C., Song, S.J., Kim, J.Y., Akita, K., Tickle, C., and Jung, H.S. (2004). Interactions between FGF and Wnt signals and Tbx3 gene expression in mammary gland initiation in mouse embryos. *Journal of anatomy* 205, 1-13.
- Eckhoute, J., Keeton, E.K., Lupien, M., Krum, S.A., Carroll, J.S., and Brown, M. (2007). Positive cross-regulatory loop ties GATA-3 to estrogen receptor alpha expression in breast cancer. *Cancer research* 67, 6477-6483.
- Efstratiadis, A., Szabolcs, M., and Klinakis, A. (2007). Notch, Myc and breast cancer. *Cell Cycle* 6, 418-429.
- Evans, M.J., and Kaufman, M.H. (1981). Establishment in culture of pluripotential cells from mouse embryos. *Nature* 292, 154-156.
- Fang, T.C., Yashiro-Ohtani, Y., Del Bianco, C., Knoblock, D.M., Blacklow, S.C., and Pear, W.S. (2007). Notch directly regulates Gata3 expression during T helper 2 cell differentiation. *Immunity* 27, 100-110.
- Feng, Y., Manka, D., Wagner, K.U., and Khan, S.A. (2007). Estrogen receptor-alpha expression in the mammary epithelium is required for ductal and alveolar morphogenesis in mice. *Proceedings of the National Academy of Sciences of the United States of America* 104, 14718-14723.
- Feng, Z., Marti, A., Jehn, B., Altermatt, H.J., Chicaiza, G., and Jaggi, R. (1995). Glucocorticoid and progesterone inhibit involution and programmed cell death in the mouse mammary gland. *The Journal of cell biology* 131, 1095-1103.
- Fiering, S., Epner, E., Robinson, K., Zhuang, Y., Telling, A., Hu, M., Martin, D.I., Enver, T., Ley, T.J., and Groudine, M. (1995). Targeted deletion of 5'HS2 of the murine beta-globin LCR reveals that it is not essential for proper regulation of the beta-globin locus. *Genes Dev* 9, 2203-2213.
- Foley, J., Dann, P., Hong, J., Cosgrove, J., Dreyer, B., Rimm, D., Dunbar, M., Philbrick, W., and Wysolmerski, J. (2001). Parathyroid hormone-related protein maintains mammary epithelial fate and triggers nipple skin differentiation during embryonic breast development. *Development (Cambridge, England)* 128, 513-525.
- Funaki, M., Katagiri, H., Inukai, K., Kikuchi, M., and Asano, T. (2000). Structure and function of phosphatidylinositol-3,4 kinase. *Cellular signalling* 12, 135-142.
- Gallahan, D., Jhappan, C., Robinson, G., Hennighausen, L., Sharp, R., Kordon, E., Callahan, R., Merlino, G., and Smith, G.H. (1996). Expression of a truncated Int3 gene in developing secretory mammary epithelium specifically retards lobular differentiation resulting in tumorigenesis. *Cancer research* 56, 1775-1785.
- Glaser, S., Anastassiadis, K., and Stewart, A.F. (2005). Current issues in mouse genome engineering. *Nature genetics* 37, 1187-1193.
- Godin, I., and Cumano, A. (2002). The hare and the tortoise: an embryonic haematopoietic race. *Nat Rev Immunol* 2, 593-604.
- Godin, I., Dieterlen-Lievre, F., and Cumano, A. (1995). Emergence of multipotent hemopoietic cells in the yolk sac and paraaortic splanchnopleura in mouse embryos,

- beginning at 8.5 days postcoitus. *Proceedings of the National Academy of Sciences of the United States of America* *92*, 773-777.
- Golonzhka, O., Leid, M., Indra, G., and Indra, A.K. (2007). Expression of COUP-TF-interacting protein 2 (CTIP2) in mouse skin during development and in adulthood. *Gene Expr Patterns* *7*, 754-760.
- Gordon, K.E., Binas, B., Chapman, R.S., Kurian, K.M., Clarkson, R.W., Clark, A.J., Lane, E.B., and Watson, C.J. (2000). A novel cell culture model for studying differentiation and apoptosis in the mouse mammary gland. *Breast Cancer Res* *2*, 222-235.
- Gossen, M., and Bujard, H. (1992). Tight control of gene expression in mammalian cells by tetracycline-responsive promoters. *Proceedings of the National Academy of Sciences of the United States of America* *89*, 5547-5551.
- Gunnarsen, J.M., Augustine, C., Spirkoska, V., Kim, M., Brown, M., and Tan, S.S. (2002). Global analysis of gene expression patterns in developing mouse neocortex using serial analysis of gene expression. *Molecular and cellular neurosciences* *19*, 560-573.
- Hameyer, D., Loonstra, A., Eshkind, L., Schmitt, S., Antunes, C., Groen, A., Bindels, E., Jonkers, J., Krimpenfort, P., Meuwissen, R., *et al.* (2007). Toxicity of ligand-dependent Cre recombinases and generation of a conditional Cre deleter mouse allowing mosaic recombination in peripheral tissues. *Physiol Genomics* *31*, 32-41.
- Hanks, M., Wurst, W., Anson-Cartwright, L., Auerbach, A.B., and Joyner, A.L. (1995). Rescue of the En-1 mutant phenotype by replacement of En-1 with En-2. *Science (New York, NY)* *269*, 679-682.
- Hardy, R.R., and Hayakawa, K. (2001). B cell development pathways. *Annual review of immunology* *19*, 595-621.
- Harris, J., Stanford, P.M., Sutherland, K., Oakes, S.R., Naylor, M.J., Robertson, F.G., Blazek, K.D., Kazlauskas, M., Hilton, H.N., Wittlin, S., *et al.* (2006). *Socs2* and *elf5* mediate prolactin-induced mammary gland development. *Mol Endocrinol* *20*, 1177-1187.
- Hatsell, S.J., and Cowin, P. (2006). Gli3-mediated repression of Hedgehog targets is required for normal mammary development. *Development (Cambridge, England)* *133*, 3661-3670.
- Heinzel, T., Lavinsky, R.M., Mullen, T.M., Soderstrom, M., Laherty, C.D., Torchia, J., Yang, W.M., Brard, G., Ngo, S.D., Davie, J.R., *et al.* (1997). A complex containing N-CoR, mSin3 and histone deacetylase mediates transcriptional repression. *Nature* *387*, 43-48.
- Hennighausen, L., and Robinson, G.W. (1998). Think globally, act locally: the making of a mouse mammary gland. *Genes Dev* *12*, 449-455.
- Hennighausen, L., and Robinson, G.W. (2005). Information networks in the mammary gland. *Nature reviews* *6*, 715-725.
- Hennighausen, L., and Robinson, G.W. (2008). Interpretation of cytokine signaling through the transcription factors STAT5A and STAT5B. *Genes Dev* *22*, 711-721.
- Hens, J.R., and Wysolmerski, J.J. (2005). Key stages of mammary gland development: molecular mechanisms involved in the formation of the embryonic mammary gland. *Breast Cancer Res* *7*, 220-224.

- Ho, I.C., and Pai, S.Y. (2007). GATA-3 - not just for Th2 cells anymore. *Cellular & molecular immunology* 4, 15-29.
- Hooper, J.E., and Scott, M.P. (2005). Communicating with Hedgehogs. *Nature reviews* 6, 306-317.
- Horseman, N.D., Zhao, W., Montecino-Rodriguez, E., Tanaka, M., Nakashima, K., Engle, S.J., Smith, F., Markoff, E., and Dorshkind, K. (1997). Defective mammopoiesis, but normal hematopoiesis, in mice with a targeted disruption of the prolactin gene. *The EMBO journal* 16, 6926-6935.
- Howard, B., Panchal, H., McCarthy, A., and Ashworth, A. (2005). Identification of the scaramanga gene implicates Neuregulin3 in mammary gland specification. *Genes Dev* 19, 2078-2090.
- Hu, C., Dievart, A., Lupien, M., Calvo, E., Tremblay, G., and Jolicoeur, P. (2006). Overexpression of activated murine Notch1 and Notch3 in transgenic mice blocks mammary gland development and induces mammary tumors. *The American journal of pathology* 168, 973-990.
- Humphreys, R.C., Bierie, B., Zhao, L., Raz, R., Levy, D., and Hennighausen, L. (2002). Deletion of Stat3 blocks mammary gland involution and extends functional competence of the secretory epithelium in the absence of lactogenic stimuli. *Endocrinology* 143, 3641-3650.
- Humphreys, R.C., Krajewska, M., Krnacik, S., Jaeger, R., Weiher, H., Krajewski, S., Reed, J.C., and Rosen, J.M. (1996). Apoptosis in the terminal endbud of the murine mammary gland: a mechanism of ductal morphogenesis. *Development (Cambridge, England)* 122, 4013-4022.
- Jhappan, C., Gallahan, D., Stahle, C., Chu, E., Smith, G.H., Merlino, G., and Callahan, R. (1992). Expression of an activated Notch-related int-3 transgene interferes with cell differentiation and induces neoplastic transformation in mammary and salivary glands. *Genes Dev* 6, 345-355.
- Johnson, R.L., and Tabin, C.J. (1997). Molecular models for vertebrate limb development. *Cell* 90, 979-990.
- Jonkers, J., and Berns, A. (1996). Retroviral insertional mutagenesis as a strategy to identify cancer genes. *Biochimica et biophysica acta* 1287, 29-57.
- Kamakura, S., Oishi, K., Yoshimatsu, T., Nakafuku, M., Masuyama, N., and Gotoh, Y. (2004). Hes binding to STAT3 mediates crosstalk between Notch and JAK-STAT signalling. *Nat Cell Biol* 6, 547-554.
- Kamimura, K., Ohi, H., Kubota, T., Okazuka, K., Yoshikai, Y., Wakabayashi, Y., Aoyagi, Y., Mishima, Y., and Kominami, R. (2007). Haploinsufficiency of Bcl11b for suppression of lymphomagenesis and thymocyte development. *Biochemical and biophysical research communications* 355, 538-542.
- Katso, R., Okkenhaug, K., Ahmadi, K., White, S., Timms, J., and Waterfield, M.D. (2001). Cellular function of phosphoinositide 3-kinases: implications for development, homeostasis, and cancer. *Annual review of cell and developmental biology* 17, 615-675.
- Kaufman, C.K., Zhou, P., Pasolli, H.A., Rendl, M., Bolotin, D., Lim, K.C., Dai, X., Alegre, M.L., and Fuchs, E. (2003). GATA-3: an unexpected regulator of cell lineage determination in skin. *Genes Dev* 17, 2108-2122.

- Kazansky, A.V., Raught, B., Lindsey, S.M., Wang, Y.F., and Rosen, J.M. (1995). Regulation of mammary gland factor/Stat5a during mammary gland development. *Mol Endocrinol* 9, 1598-1609.
- Kerr, J.F., Wyllie, A.H., and Currie, A.R. (1972). Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics. *British journal of cancer* 26, 239-257.
- Khaled, W.T., Read, E.K., Nicholson, S.E., Baxter, F.O., Brennan, A.J., Came, P.J., Sprigg, N., McKenzie, A.N., and Watson, C.J. (2007). The IL-4/IL-13/Stat6 signalling pathway promotes luminal mammary epithelial cell development. *Development (Cambridge, England)* 134, 2739-2750.
- Kleinsmith, L.J., and Pierce, G.B., Jr. (1964). Multipotentiality of Single Embryonal Carcinoma Cells. *Cancer research* 24, 1544-1551.
- Koller, B.H., and Smithies, O. (1989). Inactivating the beta 2-microglobulin locus in mouse embryonic stem cells by homologous recombination. *Proceedings of the National Academy of Sciences of the United States of America* 86, 8932-8935.
- Korach, K.S. (2000). Estrogen receptor knock-out mice: molecular and endocrine phenotypes. *Journal of the Society for Gynecologic Investigation* 7, S16-17.
- Kordon, E.C., and Smith, G.H. (1998). An entire functional mammary gland may comprise the progeny from a single cell. *Development (Cambridge, England)* 125, 1921-1930.
- Kouros-Mehr, H., Kim, J.W., Bechis, S.K., and Werb, Z. (2008). GATA-3 and the regulation of the mammary luminal cell fate. *Current opinion in cell biology* 20, 164-170.
- Kouros-Mehr, H., Slorach, E.M., Sternlicht, M.D., and Werb, Z. (2006). GATA-3 maintains the differentiation of the luminal cell fate in the mammary gland. *Cell* 127, 1041-1055.
- Kratochwil, K., and Schwartz, P. (1976). Tissue interaction in androgen response of embryonic mammary rudiment of mouse: identification of target tissue for testosterone. *Proceedings of the National Academy of Sciences of the United States of America* 73, 4041-4044.
- Kreitzer, A.C., and Regehr, W.G. (2001). Retrograde inhibition of presynaptic calcium influx by endogenous cannabinoids at excitatory synapses onto Purkinje cells. *Neuron* 29, 717-727.
- Kritikou, E.A., Sharkey, A., Abell, K., Came, P.J., Anderson, E., Clarkson, R.W., and Watson, C.J. (2003). A dual, non-redundant, role for LIF as a regulator of development and STAT3-mediated cell death in mammary gland. *Development (Cambridge, England)* 130, 3459-3468.
- Kubo, M., Hanada, T., and Yoshimura, A. (2003). Suppressors of cytokine signaling and immunity. *Nat Immunol* 4, 1169-1176.
- Lafontaine, D., and Tollervey, D. (1996). One-step PCR mediated strategy for the construction of conditionally expressed and epitope tagged yeast proteins. *Nucleic Acids Res* 24, 3469-3471.
- Lakso, M., Sauer, B., Mosinger, B., Jr., Lee, E.J., Manning, R.W., Yu, S.H., Mulder, K.L., and Westphal, H. (1992). Targeted oncogene activation by site-specific recombination in transgenic mice. *Proceedings of the National Academy of Sciences of the United States of America* 89, 6232-6236.

- LaMarca, H.L., and Rosen, J.M. (2007). Estrogen regulation of mammary gland development and breast cancer: amphiregulin takes center stage. *Breast Cancer Res* 9, 304.
- Lander, E.S., Linton, L.M., Birren, B., Nusbaum, C., Zody, M.C., Baldwin, J., Devon, K., Dewar, K., Doyle, M., FitzHugh, W., *et al.* (2001). Initial sequencing and analysis of the human genome. *Nature* 409, 860-921.
- Lee, E.C., Yu, D., Martinez de Velasco, J., Tessarollo, L., Swing, D.A., Court, D.L., Jenkins, N.A., and Copeland, N.G. (2001). A highly efficient Escherichia coli-based chromosome engineering system adapted for recombinogenic targeting and subcloning of BAC DNA. *Genomics* 73, 56-65.
- Leid, M., Ishmael, J.E., Avram, D., Shepherd, D., Fraulob, V., and Dolle, P. (2004). CTIP1 and CTIP2 are differentially expressed during mouse embryogenesis. *Gene Expr Patterns* 4, 733-739.
- Li, M., Liu, X., Robinson, G., Bar-Peled, U., Wagner, K.U., Young, W.S., Hennighausen, L., and Furth, P.A. (1997). Mammary-derived signals activate programmed cell death during the first stage of mammary gland involution. *Proceedings of the National Academy of Sciences of the United States of America* 94, 3425-3430.
- Lim, K.C., Lakshmanan, G., Crawford, S.E., Gu, Y., Grosveld, F., and Engel, J.D. (2000). Gata3 loss leads to embryonic lethality due to noradrenaline deficiency of the sympathetic nervous system. *Nature genetics* 25, 209-212.
- Lin, T.P. (1966). Microinjection of mouse eggs. *Science (New York, NY)* 151, 333-337.
- Lindeman, G.J., and Visvader, J.E. (2006). Shedding light on mammary stem cells and tumorigenesis. *Cell Cycle* 5, 671-672.
- Lindeman, G.J., Wittlin, S., Lada, H., Naylor, M.J., Santamaria, M., Zhang, J.G., Starr, R., Hilton, D.J., Alexander, W.S., Ormandy, C.J., *et al.* (2001). SOCS1 deficiency results in accelerated mammary gland development and rescues lactation in prolactin receptor-deficient mice. *Genes Dev* 15, 1631-1636.
- Liu, H., Ippolito, G.C., Wall, J.K., Niu, T., Probst, L., Lee, B.S., Pulford, K., Banham, A.H., Stockwin, L., Shaffer, A.L., *et al.* (2006). Functional studies of BCL11A: characterization of the conserved BCL11A-XL splice variant and its interaction with BCL6 in nuclear paraspeckles of germinal center B cells. *Mol Cancer* 5, 18.
- Liu, P., Jenkins, N.A., and Copeland, N.G. (2003a). A highly efficient recombineering-based method for generating conditional knockout mutations. *Genome Res* 13, 476-484.
- Liu, P., Keller, J.R., Ortiz, M., Tessarollo, L., Rachel, R.A., Nakamura, T., Jenkins, N.A., and Copeland, N.G. (2003b). Bcl11a is essential for normal lymphoid development. *Nat Immunol* 4, 525-532.
- Liu, X., Gallego, M.I., Smith, G.H., Robinson, G.W., and Hennighausen, L. (1998). Functional rescue of Stat5a-null mammary tissue through the activation of compensating signals including Stat5b. *Cell Growth Differ* 9, 795-803.
- Liu, X., Robinson, G.W., and Hennighausen, L. (1996). Activation of Stat5a and Stat5b by tyrosine phosphorylation is tightly linked to mammary gland differentiation. *Mol Endocrinol* 10, 1496-1506.
- Liu, X., Robinson, G.W., Wagner, K.U., Garrett, L., Wynshaw-Boris, A., and Hennighausen, L. (1997). Stat5a is mandatory for adult mammary gland development and lactogenesis. *Genes Dev* 11, 179-186.

- Lloyd, R.G. (1974). The segregation of the SbcA and Rac phenotypes in an *Escherichia coli* recB mutant. *Mol Gen Genet* 134, 249-259.
- Loonstra, A., Vooijs, M., Beverloo, H.B., Allak, B.A., van Drunen, E., Kanaar, R., Berns, A., and Jonkers, J. (2001). Growth inhibition and DNA damage induced by Cre recombinase in mammalian cells. *Proceedings of the National Academy of Sciences of the United States of America* 98, 9209-9214.
- Lund, L.R., Romer, J., Thomasset, N., Solberg, H., Pyke, C., Bissell, M.J., Dano, K., and Werb, Z. (1996). Two distinct phases of apoptosis in mammary gland involution: proteinase-independent and -dependent pathways. *Development (Cambridge, England)* 122, 181-193.
- Lydon, J.P., DeMayo, F.J., Funk, C.R., Mani, S.K., Hughes, A.R., Montgomery, C.A., Jr., Shyamala, G., Conneely, O.M., and O'Malley, B.W. (1995). Mice lacking progesterone receptor exhibit pleiotropic reproductive abnormalities. *Genes Dev* 9, 2266-2278.
- Mailleux, A.A., Spencer-Dene, B., Dillon, C., Ndiaye, D., Savona-Baron, C., Itoh, N., Kato, S., Dickson, C., Thiery, J.P., and Bellusci, S. (2002). Role of FGF10/FGFR2b signaling during mammary gland development in the mouse embryo. *Development (Cambridge, England)* 129, 53-60.
- Mallepell, S., Krust, A., Chambon, P., and Briskin, C. (2006). Paracrine signaling through the epithelial estrogen receptor alpha is required for proliferation and morphogenesis in the mammary gland. *Proceedings of the National Academy of Sciences of the United States of America* 103, 2196-2201.
- Marti, A., Feng, Z., Altermatt, H.J., and Jaggi, R. (1997). Milk accumulation triggers apoptosis of mammary epithelial cells. *European journal of cell biology* 73, 158-165.
- Martin, G.R. (1981). Isolation of a pluripotent cell line from early mouse embryos cultured in medium conditioned by teratocarcinoma stem cells. *Proceedings of the National Academy of Sciences of the United States of America* 78, 7634-7638.
- Martin, G.R., and Evans, M.J. (1975). Differentiation of clonal lines of teratocarcinoma cells: formation of embryoid bodies in vitro. *Proceedings of the National Academy of Sciences of the United States of America* 72, 1441-1445.
- Menzel, S., Garner, C., Gut, I., Matsuda, F., Yamaguchi, M., Heath, S., Foglio, M., Zelenika, D., Boland, A., Rooks, H., *et al.* (2007). A QTL influencing F cell production maps to a gene encoding a zinc-finger protein on chromosome 2p15. *Nature genetics* 39, 1197-1199.
- Metzger, D., and Chambon, P. (2001). Site- and time-specific gene targeting in the mouse. *Methods (San Diego, Calif)* 24, 71-80.
- Mikkola, M.L., and Millar, S.E. (2006). The mammary bud as a skin appendage: unique and shared aspects of development. *J Mammary Gland Biol Neoplasia* 11, 187-203.
- Miyoshi, K., Meyer, B., Gruss, P., Cui, Y., Renou, J.P., Morgan, F.V., Smith, G.H., Reichenstein, M., Shani, M., Hennighausen, L., *et al.* (2002). Mammary epithelial cells are not able to undergo pregnancy-dependent differentiation in the absence of the helix-loop-helix inhibitor Id2. *Mol Endocrinol* 16, 2892-2901.
- Miyoshi, K., Shillingford, J.M., Smith, G.H., Grimm, S.L., Wagner, K.U., Oka, T., Rosen, J.M., Robinson, G.W., and Hennighausen, L. (2001). Signal transducer and activator of transcription (Stat) 5 controls the proliferation and differentiation of mammary alveolar epithelium. *The Journal of cell biology* 155, 531-542.

- Mueller, S.O., Clark, J.A., Myers, P.H., and Korach, K.S. (2002). Mammary gland development in adult mice requires epithelial and stromal estrogen receptor alpha. *Endocrinology* *143*, 2357-2365.
- Murphy, K.C. (1998). Use of bacteriophage lambda recombination functions to promote gene replacement in *Escherichia coli*. *Journal of bacteriology* *180*, 2063-2071.
- Nagy, A. (2000). Cre recombinase: the universal reagent for genome tailoring. *Genesis* *26*, 99-109.
- Nakamura, T., Yamazaki, Y., Saiki, Y., Moriyama, M., Largaespada, D.A., Jenkins, N.A., and Copeland, N.G. (2000). Evi9 encodes a novel zinc finger protein that physically interacts with BCL6, a known human B-cell proto-oncogene product. *Mol Cell Biol* *20*, 3178-3186.
- Nguyen, A.V., and Pollard, J.W. (2000). Transforming growth factor beta3 induces cell death during the first stage of mammary gland involution. *Development (Cambridge, England)* *127*, 3107-3118.
- Nie, L., Perry, S.S., Zhao, Y., Huang, J., Kincade, P.W., Farrar, M.A., and Sun, X.H. (2008). Regulation of lymphocyte development by cell-type-specific interpretation of notch signals. *Mol Cell Biol* *28*, 2078-2090.
- Nishimori, K., Young, L.J., Guo, Q., Wang, Z., Insel, T.R., and Matzuk, M.M. (1996). Oxytocin is required for nursing but is not essential for parturition or reproductive behavior. *Proceedings of the National Academy of Sciences of the United States of America* *93*, 11699-11704.
- O'Connor, M., Peifer, M., and Bender, W. (1989). Construction of large DNA segments in *Escherichia coli*. *Science (New York, NY)* *244*, 1307-1312.
- Oakes, S.R., Naylor, M.J., Asselin-Labat, M.L., Blazek, K.D., Gardiner-Garden, M., Hilton, H.N., Kazlauskas, M., Pritchard, M.A., Chodosh, L.A., Pfeffer, P.L., *et al.* (2008). The Ets transcription factor Elf5 specifies mammary alveolar cell fate. *Genes Dev* *22*, 581-586.
- Okazuka, K., Wakabayashi, Y., Kashihara, M., Inoue, J., Sato, T., Yokoyama, M., Aizawa, S., Aizawa, Y., Mishima, Y., and Kominami, R. (2005). p53 prevents maturation of T cell development to the immature CD4-CD8+ stage in *Bcl11b*^{-/-} mice. *Biochemical and biophysical research communications* *328*, 545-549.
- Orkin, S.H., and Zon, L.I. (2008). Hematopoiesis: an evolving paradigm for stem cell biology. *Cell* *132*, 631-644.
- Ormandy, C.J., Camus, A., Barra, J., Damotte, D., Lucas, B., Buteau, H., Edery, M., Brousse, N., Babinet, C., Binart, N., *et al.* (1997). Null mutation of the prolactin receptor gene produces multiple reproductive defects in the mouse. *Genes Dev* *11*, 167-178.
- Perou, C.M., Sorlie, T., Eisen, M.B., van de Rijn, M., Jeffrey, S.S., Rees, C.A., Pollack, J.R., Ross, D.T., Johnsen, H., Akslen, L.A., *et al.* (2000). Molecular portraits of human breast tumours. *Nature* *406*, 747-752.
- Pollard, J.W., and Hennighausen, L. (1994). Colony stimulating factor 1 is required for mammary gland development during pregnancy. *Proceedings of the National Academy of Sciences of the United States of America* *91*, 9312-9316.
- Przybylski, G.K., Dik, W.A., Wanzeck, J., Grabarczyk, P., Majunke, S., Martin-Subero, J.I., Siebert, R., Dolken, G., Ludwig, W.D., Verhaaf, B., *et al.* (2005). Disruption of the BCL11B gene through *inv(14)(q11.2q32.31)* results in the expression of BCL11B-

- TRDC fusion transcripts and is associated with the absence of wild-type BCL11B transcripts in T-ALL. *Leukemia* 19, 201-208.
- Pulford, K., Banham, A.H., Lyne, L., Jones, M., Ippolito, G.C., Liu, H., Tucker, P.W., Roncador, G., Lucas, E., Ashe, S., *et al.* (2006). The BCL11AXL transcription factor: its distribution in normal and malignant tissues and use as a marker for plasmacytoid dendritic cells. *Leukemia* 20, 1439-1441.
- Radtke, F., Wilson, A., and MacDonald, H.R. (2004). Notch signaling in T- and B-cell development. *Current opinion in immunology* 16, 174-179.
- Rajewsky, K., Gu, H., Kuhn, R., Betz, U.A., Muller, W., Roes, J., and Schwenk, F. (1996). Conditional gene targeting. *The Journal of clinical investigation* 98, 600-603.
- Ramirez-Solis, R., Davis, A.C., and Bradley, A. (1993). Gene targeting in embryonic stem cells. *Methods in enzymology* 225, 855-878.
- Ramirez-Solis, R., Liu, P., and Bradley, A. (1995). Chromosome engineering in mice. *Nature* 378, 720-724.
- Raouf, A., Zhao, Y., To, K., Stingl, J., Delaney, A., Barbara, M., Iscove, N., Jones, S., McKinney, S., Emerman, J., *et al.* (2008). Transcriptome analysis of the normal human mammary cell commitment and differentiation process. *Cell stem cell* 3, 109-118.
- Raught, B., Liao, W.S., and Rosen, J.M. (1995). Developmentally and hormonally regulated CCAAT/enhancer-binding protein isoforms influence beta-casein gene expression. *Mol Endocrinol* 9, 1223-1232.
- Raymond, C.S., and Soriano, P. (2007). High-efficiency FLP and PhiC31 site-specific recombination in mammalian cells. *PLoS ONE* 2, e162.
- Robertson, E., Bradley, A., Kuehn, M., and Evans, M. (1986). Germ-line transmission of genes introduced into cultured pluripotential cells by retroviral vector. *Nature* 323, 445-448.
- Robertson, G., Hirst, M., Bainbridge, M., Bilenky, M., Zhao, Y., Zeng, T., Euskirchen, G., Bernier, B., Varhol, R., Delaney, A., *et al.* (2007). Genome-wide profiles of STAT1 DNA association using chromatin immunoprecipitation and massively parallel sequencing. *Nature methods* 4, 651-657.
- Rohr, O., Lecestre, D., Chasserot-Golaz, S., Marban, C., Avram, D., Aunis, D., Leid, M., and Schaeffer, E. (2003). Recruitment of Tat to heterochromatin protein HP1 via interaction with CTIP2 inhibits human immunodeficiency virus type 1 replication in microglial cells. *Journal of virology* 77, 5415-5427.
- Rothenberg, E.V. (2007a). Cell lineage regulators in B and T cell development. *Nat Immunol* 8, 441-444.
- Rothenberg, E.V. (2007b). Regulatory factors for initial T lymphocyte lineage specification. *Current opinion in hematology* 14, 322-329.
- Russo, J., Ao, X., Grill, C., and Russo, I.H. (1999). Pattern of distribution of cells positive for estrogen receptor alpha and progesterone receptor in relation to proliferating cells in the mammary gland. *Breast cancer research and treatment* 53, 217-227.
- Saiki, Y., Yamazaki, Y., Yoshida, M., Katoh, O., and Nakamura, T. (2000). Human EVI9, a homologue of the mouse myeloid leukemia gene, is expressed in the hematopoietic progenitors and down-regulated during myeloid differentiation of HL60 cells. *Genomics* 70, 387-391.

- Sakakura, T., Kusano, I., Kusakabe, M., Inaguma, Y., and Nishizuka, Y. (1987). Biology of mammary fat pad in fetal mouse: capacity to support development of various fetal epithelia in vivo. *Development (Cambridge, England)* *100*, 421-430.
- Satokata, I., Ma, L., Ohshima, H., Bei, M., Woo, I., Nishizawa, K., Maeda, T., Takano, Y., Uchiyama, M., Heaney, S., *et al.* (2000). Msx2 deficiency in mice causes pleiotropic defects in bone growth and ectodermal organ formation. *Nature genetics* *24*, 391-395.
- Satterwhite, E., Sonoki, T., Willis, T.G., Harder, L., Nowak, R., Arriola, E.L., Liu, H., Price, H.P., Gesk, S., Steinemann, D., *et al.* (2001). The BCL11 gene family: involvement of BCL11A in lymphoid malignancies. *Blood* *98*, 3413-3420.
- Sauer, B., and Henderson, N. (1988). Site-specific DNA recombination in mammalian cells by the Cre recombinase of bacteriophage P1. *Proceedings of the National Academy of Sciences of the United States of America* *85*, 5166-5170.
- Schlake, T., and Bode, J. (1994). Use of mutated FLP recognition target (FRT) sites for the exchange of expression cassettes at defined chromosomal loci. *Biochemistry* *33*, 12746-12751.
- Schorr, K., Li, M., Krajewski, S., Reed, J.C., and Furth, P.A. (1999). Bcl-2 gene family and related proteins in mammary gland involution and breast cancer. *J Mammary Gland Biol Neoplasia* *4*, 153-164.
- Seagroves, T.N., Krnacik, S., Raught, B., Gay, J., Burgess-Beusse, B., Darlington, G.J., and Rosen, J.M. (1998). C/EBPbeta, but not C/EBPalpha, is essential for ductal morphogenesis, lobuloalveolar proliferation, and functional differentiation in the mouse mammary gland. *Genes Dev* *12*, 1917-1928.
- Selbert, S., Bentley, D.J., Melton, D.W., Rannie, D., Lourenco, P., Watson, C.J., and Clarke, A.R. (1998). Efficient BLG-Cre mediated gene deletion in the mammary gland. *Transgenic Res* *7*, 387-396.
- Senawong, T., Peterson, V.J., Avram, D., Shepherd, D.M., Frye, R.A., Minucci, S., and Leid, M. (2003). Involvement of the histone deacetylase SIRT1 in chicken ovalbumin upstream promoter transcription factor (COUP-TF)-interacting protein 2-mediated transcriptional repression. *J Biol Chem* *278*, 43041-43050.
- Senawong, T., Peterson, V.J., and Leid, M. (2005). BCL11A-dependent recruitment of SIRT1 to a promoter template in mammalian cells results in histone deacetylation and transcriptional repression. *Arch Biochem Biophys* *434*, 316-325.
- Shackleton, M., Vaillant, F., Simpson, K.J., Stingl, J., Smyth, G.K., Asselin-Labat, M.L., Wu, L., Lindeman, G.J., and Visvader, J.E. (2006). Generation of a functional mammary gland from a single stem cell. *Nature* *439*, 84-88.
- Shillingford, J.M., Miyoshi, K., Robinson, G.W., Bierie, B., Cao, Y., Karin, M., and Hennighausen, L. (2003). Proteotyping of mammary tissue from transgenic and gene knockout mice with immunohistochemical markers: a tool to define developmental lesions. *J Histochem Cytochem* *51*, 555-565.
- Shinbo, T., Matsuki, A., Matsumoto, Y., Kosugi, S., Takahashi, Y., Niwa, O., and Kominami, R. (1999). Allelic loss mapping and physical delineation of a region harboring a putative thymic lymphoma suppressor gene on mouse chromosome 12. *Oncogene* *18*, 4131-4136.
- Silver, L.M. (1985). *Mouse Genetics, Concepts and Applications* (Oxford University Press).

- Singh, H., Medina, K.L., and Pongubala, J.M. (2005). Contingent gene regulatory networks and B cell fate specification. *Proceedings of the National Academy of Sciences of the United States of America* *102*, 4949-4953.
- Sleeman, K.E., Kendrick, H., Robertson, D., Isacke, C.M., Ashworth, A., and Smalley, M.J. (2007). Dissociation of estrogen receptor expression and in vivo stem cell activity in the mammary gland. *The Journal of cell biology* *176*, 19-26.
- Smalley, M.J., Titley, J., and O'Hare, M.J. (1998). Clonal characterization of mouse mammary luminal epithelial and myoepithelial cells separated by fluorescence-activated cell sorting. *In vitro cellular & developmental biology* *34*, 711-721.
- Smith, G.H. (1996). Experimental mammary epithelial morphogenesis in an in vivo model: evidence for distinct cellular progenitors of the ductal and lobular phenotype. *Breast cancer research and treatment* *39*, 21-31.
- Smith, G.H. (2005). Label-retaining epithelial cells in mouse mammary gland divide asymmetrically and retain their template DNA strands. *Development (Cambridge, England)* *132*, 681-687.
- Smith, G.H., and Boulanger, C.A. (2003). Mammary epithelial stem cells: transplantation and self-renewal analysis. *Cell proliferation* *36 Suppl 1*, 3-15.
- Smith, G.H., Gallahan, D., Diella, F., Jhappan, C., Merlino, G., and Callahan, R. (1995). Constitutive expression of a truncated INT3 gene in mouse mammary epithelium impairs differentiation and functional development. *Cell Growth Differ* *6*, 563-577.
- Smith, G.H., and Medina, D. (1988). A morphologically distinct candidate for an epithelial stem cell in mouse mammary gland. *Journal of cell science* *90 (Pt 1)*, 173-183.
- Smithies, O., Gregg, R.G., Boggs, S.S., Koralewski, M.A., and Kucherlapati, R.S. (1985). Insertion of DNA sequences into the human chromosomal beta-globin locus by homologous recombination. *Nature* *317*, 230-234.
- Snell, G.D., and Reed, S. (1993). William Ernest Castle, pioneer mammalian geneticist. *Genetics* *133*, 751-753.
- Solter, D., Skreb, N., and Damjanov, I. (1970). Extrauterine growth of mouse egg-cylinders results in malignant teratoma. *Nature* *227*, 503-504.
- Sorlie, T., Perou, C.M., Tibshirani, R., Aas, T., Geisler, S., Johnsen, H., Hastie, T., Eisen, M.B., van de Rijn, M., Jeffrey, S.S., *et al.* (2001). Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. *Proceedings of the National Academy of Sciences of the United States of America* *98*, 10869-10874.
- Spangrude, G.J., Heimfeld, S., and Weissman, I.L. (1988). Purification and characterization of mouse hematopoietic stem cells. *Science (New York, NY)* *241*, 58-62.
- Sternlicht, M.D. (2006). Key stages in mammary gland development: the cues that regulate ductal branching morphogenesis. *Breast Cancer Res* *8*, 201.
- Stingl, J., and Caldas, C. (2007). Molecular heterogeneity of breast carcinomas and the cancer stem cell hypothesis. *Nat Rev Cancer* *7*, 791-799.
- Stingl, J., Eaves, C.J., Zandieh, I., and Emerman, J.T. (2001). Characterization of bipotent mammary epithelial progenitor cells in normal adult human breast tissue. *Breast cancer research and treatment* *67*, 93-109.

- Stingl, J., Eirew, P., Ricketson, I., Shackleton, M., Vaillant, F., Choi, D., Li, H.I., and Eaves, C.J. (2006). Purification and unique properties of mammary epithelial stem cells. *Nature* 439, 993-997.
- Stylianou, S., Clarke, R.B., and Brennan, K. (2006). Aberrant activation of notch signaling in human breast cancer. *Cancer research* 66, 1517-1525.
- Suzuki, M., Gerstein, M., and Yagi, N. (1994). Stereochemical basis of DNA recognition by Zn fingers. *Nucleic Acids Res* 22, 3397-3405.
- Taddei, I., Deugnier, M.A., Faraldo, M.M., Petit, V., Bouvard, D., Medina, D., Fassler, R., Thiery, J.P., and Glukhova, M.A. (2008). Beta1 integrin deletion from the basal compartment of the mammary epithelium affects stem cells. *Nat Cell Biol* 10, 716-722.
- Teglund, S., McKay, C., Schuetz, E., van Deursen, J.M., Stravopodis, D., Wang, D., Brown, M., Bodner, S., Grosveld, G., and Ihle, J.N. (1998). Stat5a and Stat5b proteins have essential and nonessential, or redundant, roles in cytokine responses. *Cell* 93, 841-850.
- Thangaraju, M., Rudelius, M., Bierie, B., Raffeld, M., Sharan, S., Hennighausen, L., Huang, A.M., and Sterneck, E. (2005). C/EBPdelta is a crucial regulator of pro-apoptotic gene expression during mammary gland involution. *Development (Cambridge, England)* 132, 4675-4685.
- Thomas, K.R., and Capecchi, M.R. (1986). Introduction of homologous DNA sequences into mammalian cells induces mutations in the cognate gene. *Nature* 324, 34-38.
- Thomas, K.R., Folger, K.R., and Capecchi, M.R. (1986). High frequency targeting of genes to specific sites in the mammalian genome. *Cell* 44, 419-428.
- Timmons, L., Becker, J., Barthmaier, P., Fyrberg, C., Shearn, A., and Fyrberg, E. (1997). Green fluorescent protein/beta-galactosidase double reporters for visualizing *Drosophila* gene expression patterns. *Developmental genetics* 20, 338-347.
- Tonner, E., Barber, M.C., Allan, G.J., Beattie, J., Webster, J., Whitelaw, C.B., and Flint, D.J. (2002). Insulin-like growth factor binding protein-5 (IGFBP-5) induces premature cell death in the mammary glands of transgenic mice. *Development (Cambridge, England)* 129, 4547-4557.
- Topper, Y.J., and Freeman, C.S. (1980). Multiple hormone interactions in the developmental biology of the mammary gland. *Physiological reviews* 60, 1049-1106.
- Trainor, P.A., Zhou, S.X., Parameswaran, M., Quinlan, G.A., Gordon, M., Sturm, K., and Tam, P.P. (1999). Application of lacZ transgenic mice to cell lineage studies. *Methods in molecular biology (Clifton, NJ)* 97, 183-200.
- Udy, G.B., Towers, R.P., Snell, R.G., Wilkins, R.J., Park, S.H., Ram, P.A., Waxman, D.J., and Davey, H.W. (1997). Requirement of STAT5b for sexual dimorphism of body growth rates and liver gene expression. *Proceedings of the National Academy of Sciences of the United States of America* 94, 7239-7244.
- Vaillant, F., Asselin-Labat, M.L., Shackleton, M., Lindeman, G.J., and Visvader, J.E. (2007). The emerging picture of the mouse mammary stem cell. *Stem cell reviews* 3, 114-123.
- van Deursen, J., and Wieringa, B. (1992). Targeting of the creatine kinase M gene in embryonic stem cells using isogenic and nonisogenic vectors. *Nucleic Acids Res* 20, 3815-3820.
- van Genderen, C., Okamura, R.M., Farinas, I., Quo, R.G., Parslow, T.G., Bruhn, L., and Grosschedl, R. (1994). Development of several organs that require inductive epithelial-

- mesenchymal interactions is impaired in LEF-1-deficient mice. *Genes Dev* 8, 2691-2703.
- Veltmaat, J.M., Mailleux, A.A., Thiery, J.P., and Bellusci, S. (2003). Mouse embryonic mammaryogenesis as a model for the molecular regulation of pattern formation. *Differentiation; research in biological diversity* 71, 1-17.
- Veltmaat, J.M., Van Veelen, W., Thiery, J.P., and Bellusci, S. (2004). Identification of the mammary line in mouse by Wnt10b expression. *Dev Dyn* 229, 349-356.
- Venter, J.C., Adams, M.D., Myers, E.W., Li, P.W., Mural, R.J., Sutton, G.G., Smith, H.O., Yandell, M., Evans, C.A., Holt, R.A., *et al.* (2001). The sequence of the human genome. *Science (New York, NY)* 291, 1304-1351.
- Visvader, J.E., and Lindeman, G.J. (2006). Mammary stem cells and mammaryogenesis. *Cancer research* 66, 9798-9801.
- Wagner, E.F., Stewart, T.A., and Mintz, B. (1981a). The human beta-globin gene and a functional viral thymidine kinase gene in developing mice. *Proceedings of the National Academy of Sciences of the United States of America* 78, 5016-5020.
- Wagner, K.U., Young, W.S., 3rd, Liu, X., Ginns, E.I., Li, M., Furth, P.A., and Hennighausen, L. (1997). Oxytocin and milk removal are required for post-partum mammary-gland development. *Genes and function* 1, 233-244.
- Wagner, T.E., Hoppe, P.C., Jollick, J.D., Scholl, D.R., Hodinka, R.L., and Gault, J.B. (1981b). Microinjection of a rabbit beta-globin gene into zygotes and its subsequent expression in adult mice and their offspring. *Proceedings of the National Academy of Sciences of the United States of America* 78, 6376-6380.
- Wakabayashi, Y., Inoue, J., Takahashi, Y., Matsuki, A., Kosugi-Okano, H., Shinbo, T., Mishima, Y., Niwa, O., and Kominami, R. (2003a). Homozygous deletions and point mutations of the *Rit1/Bcl11b* gene in gamma-ray induced mouse thymic lymphomas. *Biochemical and biophysical research communications* 301, 598-603.
- Wakabayashi, Y., Watanabe, H., Inoue, J., Takeda, N., Sakata, J., Mishima, Y., Hitomi, J., Yamamoto, T., Utsuyama, M., Niwa, O., *et al.* (2003b). *Bcl11b* is required for differentiation and survival of alphabeta T lymphocytes. *Nat Immunol* 4, 533-539.
- Walton, K.D., Wagner, K.U., Rucker, E.B., 3rd, Shillingford, J.M., Miyoshi, K., and Hennighausen, L. (2001). Conditional deletion of the *bcl-x* gene from mouse mammary epithelium results in accelerated apoptosis during involution but does not compromise cell function during lactation. *Mechanisms of development* 109, 281-293.
- Wang, J., Sarov, M., Rientjes, J., Fu, J., Hollak, H., Kranz, H., Xie, W., Stewart, A.F., and Zhang, Y. (2006). An improved recombineering approach by adding *RecA* to lambda Red recombination. *Molecular biotechnology* 32, 43-53.
- Warren, A.J., Colledge, W.H., Carlton, M.B., Evans, M.J., Smith, A.J., and Rabbitts, T.H. (1994). The oncogenic cysteine-rich LIM domain protein *rbtn2* is essential for erythroid development. *Cell* 78, 45-57.
- Waterston, R.H., Lindblad-Toh, K., Birney, E., Rogers, J., Abril, J.F., Agarwal, P., Agarwala, R., Ainscough, R., Alexandersson, M., An, P., *et al.* (2002). Initial sequencing and comparative analysis of the mouse genome. *Nature* 420, 520-562.
- Watson, C.J. (2006a). Involution: apoptosis and tissue remodelling that convert the mammary gland from milk factory to a quiescent organ. *Breast Cancer Res* 8, 203.
- Watson, C.J. (2006b). Post-lactational mammary gland regression: molecular basis and implications for breast cancer. *Expert Reviews in Molecular Medicine* 8, 1.

- Watson, C.J., and Khaled, W.T. (2008). Mammary development in the embryo and adult: a journey of morphogenesis and commitment. *Development (Cambridge, England)* 135, 995-1003.
- Welm, B.E., Tepera, S.B., Venezia, T., Graubert, T.A., Rosen, J.M., and Goodell, M.A. (2002). Sca-1(pos) cells in the mouse mammary gland represent an enriched progenitor cell population. *Developmental biology* 245, 42-56.
- Weniger, M.A., Pulford, K., Gesk, S., Ehrlich, S., Banham, A.H., Lyne, L., Martin-Subero, J.I., Siebert, R., Dyer, M.J., Moller, P., *et al.* (2006). Gains of the proto-oncogene BCL11A and nuclear accumulation of BCL11A(XL) protein are frequent in primary mediastinal B-cell lymphoma. *Leukemia* 20, 1880-1882.
- Wood, L.D., Parsons, D.W., Jones, S., Lin, J., Sjoblom, T., Leary, R.J., Shen, D., Boca, S.M., Barber, T., Ptak, J., *et al.* (2007). The genomic landscapes of human breast and colorectal cancers. *Science (New York, NY)* 318, 1108-1113.
- Wormald, S., and Hilton, D.J. (2004). Inhibitors of cytokine signal transduction. *J Biol Chem* 279, 821-824.
- Wysolmerski, J.J., Philbrick, W.M., Dunbar, M.E., Lanske, B., Kronenberg, H., and Broadus, A.E. (1998). Rescue of the parathyroid hormone-related protein knockout mouse demonstrates that parathyroid hormone-related protein is essential for mammary gland development. *Development (Cambridge, England)* 125, 1285-1294.
- Yamaguchi, N., Oyama, T., Ito, E., Satoh, H., Azuma, S., Hayashi, M., Shimizu, K., Honma, R., Yanagisawa, Y., Nishikawa, A., *et al.* (2008). NOTCH3 signaling pathway plays crucial roles in the proliferation of ErbB2-negative human breast cancer cells. *Cancer research* 68, 1881-1888.
- Young, W.S., 3rd, Shepard, E., Amico, J., Hennighausen, L., Wagner, K.U., LaMarca, M.E., McKinney, C., and Ginns, E.I. (1996). Deficiency in mouse oxytocin prevents milk ejection, but not fertility or parturition. *Journal of neuroendocrinology* 8, 847-853.
- Yu, D., Ellis, H.M., Lee, E.C., Jenkins, N.A., Copeland, N.G., and Court, D.L. (2000). An efficient recombination system for chromosome engineering in *Escherichia coli*. *Proceedings of the National Academy of Sciences of the United States of America* 97, 5978-5983.
- Zhang, Y., Buchholz, F., Muyrers, J.P., and Stewart, A.F. (1998). A new logic for DNA engineering using recombination in *Escherichia coli*. *Nature genetics* 20, 123-128.
- Zheng, W., and Flavell, R.A. (1997). The transcription factor GATA-3 is necessary and sufficient for Th2 cytokine gene expression in CD4 T cells. *Cell* 89, 587-596.
- Zhou, J., Chehab, R., Tkalcevic, J., Naylor, M.J., Harris, J., Wilson, T.J., Tsao, S., Tellis, I., Zavarsek, S., Xu, D., *et al.* (2005). Elf5 is essential for early embryogenesis and mammary gland development during pregnancy and lactation. *The EMBO journal* 24, 635-644.