

Biochemical investigations of *Plasmodium falciparum* erythrocyte invasion pathways using recombinant merozoite surface proteins produced in a mammalian expression system

Madushi Wanaguru

Downing College, University of Cambridge

A dissertation submitted for the degree of

Doctor of Philosophy

September 2012

For Amma

Declaration

This dissertation is the result of my own work and includes nothing which is the outcome of work done in collaboration except where specifically indicated in the text. This dissertation does not exceed the word limit set by the Biology Degree Committee.

Summary

The invasion of erythrocytes by *Plasmodium falciparum* merozoites is mediated by low-affinity, extracellular interactions between surface proteins on the host and parasite cells. Identification of these interactions *in vitro* is technically challenging; not only are *P. falciparum* proteins hard to express recombinantly in heterologous systems, but detecting their transient interactions with erythrocyte receptors is difficult using traditional biochemical methods. Two significant advances in this area of research have recently been made in our laboratory. Firstly, a strategy has been optimised for the production of soluble full-length ectodomains of *P. falciparum* merozoite surface proteins, in a mammalian expression system. Secondly, an ELISA-based method, AVEXIS (avidity-based extracellular interaction screening) has been developed which enables low-affinity interactions between proteins to be identified in a high-throughput manner. The AVEXIS assay is, however, limited for use with soluble proteins and as such cannot be used for the analysis of multi-pass membrane proteins. In my PhD, I developed a flow-cytometry based, high-throughput assay for identifying low-affinity interactions between recombinant *P. falciparum* proteins and receptors displayed on the surface of cells. I used this assay in combination with other biochemical and biophysical methods such as AVEXIS and surface plasmon resonance to investigate selected aspects/interactions of erythrocyte invasion in three projects.

P. falciparum EBA175 is the known ligand of the erythrocyte receptor Glycophorin A, which carries the antigenic determinants of the human MN blood group system. In my first project, I showed that the mammalian-expressed, full-length ectodomain of EBA175 (*PfEBA175* FL) is functionally similar to native EBA175 isolated from parasite cultures and that it binds MN Glycophorin A with a slightly higher affinity than the MM form. I also found that *PfEBA175*

RII, a truncated derivative containing only the extracellular region known to be essential for interacting with erythrocytes, binds to native Glycophorin A with a one order of magnitude lower affinity than *PfEBA175* FL, suggesting some role played by regions outside of RII to facilitate binding.

The closest relatives of *P. falciparum* are found amongst the *Laverania* family of great ape parasites, which appear to be strictly host-specific in their natural environments. In my second project, I investigated the potential contributions of EBA175-Glycophorin A and another parasite ligand-host receptor interaction, RH5-Basigin (BSG), towards the determination of host-specificity in *Laverania*. I only observed a relatively small difference in the affinities for native human Glycophorin A, between *PfEBA175* and orthologues of EBA175 from two chimpanzee parasites *P. reichenowi* and *P. billcollinsi*, which suggests that this interaction may not play a significant role in the determination of host-specificity in *Laverania*. On the other hand, I saw clear host-selectivity in the recognition of BSG orthologues from human, chimpanzee and gorilla, by *P. falciparum* RH5 and identified residues that confer this specificity by generating and testing site-directed mutants of the BSG orthologues.

Although multi-pass receptors are fairly abundant on the erythrocyte surface, no such receptor has yet been conclusively identified to play a role in erythrocyte invasion by *P. falciparum*. To potentially identify such a novel interaction, I screened a library of mammalian-expressed *P. falciparum* proteins against a panel of recombinant erythrocyte multi-pass receptors, in my third project. I also tested the library of merozoite proteins for binding to human erythrocytes and to a large panel of synthetic carbohydrates. A number of putative interactions were identified in the screens and are awaiting characterisation.

Acknowledgements

I am most indebted to Gavin Wright, for the opportunity to work in his lab and for the thorough and patient mentorship throughout the course of my PhD. I am also very grateful to Julian Rayner for his advice and encouragement.

I wish to thank Cecile Crosnier and Josefin Bartholdson for their kind help and advice, Zenon Zenonos for the anti-Basigin monoclonals and Leyla Bustamante for the human erythrocytes. I am also grateful to Bee Ng and William Cheng for flow cytometry training and Gareth Powell for confocal microscopy training. I'm lucky to have worked with my colleagues in Team 30, everyone of whom have helped me in one way or another.

I'm very obliged to my friends for standing by me through the ups and downs of PhD life, in particular Jenn and Laure, who were pillars of emotional support. Nothing of course would ever have been possible without my family; my brothers, Chamitha and Miraj and most of all my father, Bandula.

Table of Contents

Declaration	iii
Summary	iv
Acknowledgements	vi
Table of Contents	vii
Chapter 1: General Introduction	1
1.1 Malaria is a global health problem	2
1.2 Five different species of <i>Plasmodium</i> can cause malaria in humans	3
1.3 The closest relatives of <i>P. falciparum</i> are great ape parasites of the <i>Laverania</i> family	3
1.4 <i>P. falciparum</i> has a complex life cycle	6
1.5 The invasion of human erythrocytes by <i>P. falciparum</i> merozoites is a multi-step process mediated by interactions between cell-surface proteins	8
1.6 Primary contact with erythrocytes is proposed to be mediated by proteins that constitute the outer coat of <i>P. falciparum</i> merozoites	11
1.7 Merozoite proteins of the EBL and RH families play a critical role during erythrocyte invasion	13
1.8 The micronemal protein AMA1 is essential for tight junction formation	15
1.9 Adhesins may be coupled to the molecular motor by MTRAP to allow movement of the tight junction	15
1.10 <i>P. falciparum</i> merozoite surface proteins are important candidates for a blood-stage vaccine	16
1.11 Many prokaryotic and eukaryotic expression systems have been tested for the production of <i>P. falciparum</i> proteins in recombinant form	17
1.12 Studying extracellular interactions between <i>P. falciparum</i> merozoites and human erythrocytes using <i>in vitro</i> biochemical techniques is challenging	19

1.13 Work in this laboratory: expression of a library of <i>P. falciparum</i> merozoite surface proteins and avidity-based extracellular interactions screening (AVEXIS)	21
1.14 Work described in this thesis	22
1.15 Bibliography	30
Chapter 2: Materials and methods	40
2.1 Production of recombinant membrane protein ectodomains	41
2.1.1 Design and construction of expression plasmids	41
2.1.1.1 PCR	43
2.1.1.2 Agarose gel electrophoresis	45
2.1.2 Transient transfection of expression plasmids	45
2.2 Qualitative and quantitative assessment of recombinant membrane protein ectodomains	46
2.2.1 SDS-polyacrylamide gel electrophoresis (SDS-PAGE)	46
2.2.2 Western blotting	46
2.2.3 Enzyme-linked-immunosorbent assays (ELISAs) of biotinylated membrane protein ectodomains	46
2.2.4 Normalisation of β -lactamase tagged membrane protein ectodomains	47
2.2.5 Purification of His-tagged membrane protein ectodomains	47
2.3 <i>In vitro</i> biotinylation of native Glycophorin A	49
2.4 Expression and analysis of erythrocyte multi-pass membrane proteins	49
2.4.1 Transient transfection of TrueORF plasmids	49
2.4.2 Monitoring expression of multi-pass membrane proteins by flow cytometry	51
2.4.3 Confirming expression of multi-pass membrane proteins by confocal microscopy	51
2.5 Identifying and characterising low-affinity protein-protein interactions	52
2.5.1 Cell-based binding assays with ‘prey’ protein arrays immobilised on beads	52

2.5.1.1 Pre-treatment of cells	53
2.5.2 Avidity-based extracellular interaction screen (AVEXIS)	54
2.5.3 Surface plasmon resonance (SPR)	54
2.6 Studying protein-glycan interactions	56
2.6.1 Lectin binding assay	56
2.6.2 High-throughput screen for identifying glycan-binding specificities of <i>P. falciparum</i> membrane proteins	56
2.7 Tables	57
2.8 Bibliography	62
Chapter 3: Functional validation of a mammalian-expressed EBA175 antigen and comparative analysis of its binding to MM and MN forms of human Glycophorin A	64
3.1 Introduction	65
3.1.1 Identification and characterisation of <i>P. falciparum</i> EBA175 (<i>PfEBA175</i>)	65
3.1.2 Interaction of <i>PfEBA175</i> with the erythrocyte receptor, Glycophorin A	68
3.1.3 <i>PfEBA175</i> is an important vaccine candidate	70
3.1.4 Work described in the chapter	71
3.2 Results	
3.2.1 <i>PfEBA175</i> FL was expressed as a soluble fusion protein and immunologically characterised.	72
3.2.2 Recombinant <i>PfEBA175</i> FL showed Glycophorin A-dependent binding to human erythrocytes.	76
3.2.3 Recombinant <i>PfEBA175</i> FL showed binding to native human Glycophorin A in a sialic acid-dependent manner	82
3.2.4 The full-length ectodomain of human Glycophorin A was expressed in soluble recombinant form	88
3.2.5 No binding was detected between recombinant Glycophorin A and	89

<i>Pf</i> EBA175 FL using AVEXIS	
3.2.6 No binding of recombinant Glycophorin A to <i>Pf</i> EBA175 FL was detected by SPR	92
3.2.7 The level of sialylation of recombinant Glycophorin A was increased by co-expression with sialyltransferases and a sialic acid transporter	94
3.2.8 Kinetic parameters for the interaction between native Glycophorin A and <i>Pf</i> EBA175 FL were estimated by SPR	97
3.2.9 <i>Pf</i> EBA175 RII was produced in soluble form and its interactions with native Glycophorin A was probed	110
3.3 Discussion	
3.3.1 Recombinant <i>Pf</i> EBA175 FL is functionally similar to native <i>Pf</i> EBA175 isolated from parasite cultures	116
3.3.2 The kinetic profile of the binding of <i>Pf</i> EBA175 FL to native Glycophorin A is consistent with the ‘ligand-induced dimerisation model’	118
3.3.3 The affinity of <i>Pf</i> EBA175 FL for MN Glycophorin A is x 1.3-fold higher than that for MM Glycophorin A	119
3.3.4 <i>Pf</i> EBA175 FL binds to native Glycophorin A with a ~10-fold higher affinity than <i>Pf</i> EBA175 RII	120
3.3.5 Functional activity of the extracellular domain of Glycophorin A may be dependent on factors other than sialylation	121
3.4 Conclusion	122
3.5 Bibliography	123
Chapter 4: Investigating the host-specificity of <i>Plasmodium</i> merozoite: primate erythrocyte interactions in the <i>Laverania</i> family	127
4.1 Introduction	128
4.1.1 The <i>Laverania</i> family of great ape parasites are host-specific	128
4.1.2 The molecular-basis of the <i>Laverania</i> host-specificity is poorly understood	129
4.1.3 The RH5-Basigin interaction as a possible determinant of host-specificity	130

4.1.4 Work described in this chapter	131
4.2 Results	
4.2.1 EBA175 RII orthologues from three <i>Laverania</i> species were expressed and immunologically characterised	132
4.2.2 All three EBA175 RII orthologues showed specific binding to human erythrocytes	135
4.2.3 Purified native human Glycophorin A was recognized by all three EBA175 RII orthologues in a sialic acid-dependent manner	137
4.2.4 SPR studies revealed differences in the affinities of the EBA175 RII orthologues for human Glycophorin A	142
4.2.5 RH5 orthologues from two <i>Laverania</i> species were expressed and biochemically characterised	148
4.2.6 Both RH5 orthologues showed sialic acid-independent binding to human erythrocytes	150
4.2.7 BSG orthologues from three species of primates were expressed and characterised	153
4.2.8 Differences in the binding of RH5 to the BSG orthologues were revealed by AVEXIS	158
4.2.9 Affinity measurements for the RH5-BSG interaction were obtained by SPR	162
4.2.10 Site-directed mutants of human and chimpanzee BSG proteins were generated and tested for correct folding	169
4.2.11 The affinities of BSG mutants for <i>P. falciparum</i> RH5 were compared to those of human and chimpanzee BSG	171
4.3 Discussion	177
4.3.1 The EBA175-Glycophorin A interaction may not contribute significantly to the establishment of host-specificity in <i>Laverania</i>	178
4.3.2 The RH5-BSG interaction is likely to be a significant determinant of <i>Laverania</i> host-specificity	182
4.4 Conclusion	185

4.5 Bibliography	188
Chapter 5: Development of high-throughput assays for characterising a library of <i>P. falciparum</i> merozoite surface proteins	191
5.1 Introduction	192
5.1.1 Cell surface proteins of the human erythrocyte	192
5.1.2 The role of host-cell surface multi-pass membrane proteins in the invasion of erythrocytes by <i>Plasmodium</i> merozoites	192
5.1.3 The recognition of glycan moieties on erythrocyte surface receptors by <i>Plasmodium</i> ligands	195
5.1.4 Strategies for detecting and characterising interactions between erythrocyte and <i>Plasmodium</i> merozoite surface proteins	195
5.1.5 Work described in this chapter	196
5.2 Results	199
5.2.1 40% of the <i>P. falciparum</i> merozoite surface proteins tested showed some association with human erythrocytes	199
5.2.2 An expression library of erythrocyte multi-pass proteins was compiled based on published literature and protein topology modelling	206
5.2.3 Proof-of-principle study: demonstration of the interaction between Cd200 and its receptor with one partner expressed transiently on the surface of HEK293E cells and the other immobilised on fluorescent beads	211
5.2.4 Positive control study: demonstration of the interaction between the Duffy antigen expressed recombinantly on the surface of HEK293E cells by transient transfection and PvDBP immobilised on fluorescent beads	213
5.2.5 More than 40% of <i>P. falciparum</i> merozoite surface proteins tested showed binding to untreated and mock-transfected HEK293E cells	214
5.2.6 The <i>P. falciparum</i> merozoite surface proteins were screened against 41 erythrocytic multi-pass receptors, each expressed individually on the surface of HEK293E cells	222
5.2.7 Preliminary follow-up study: recapitulation of the interaction between AARP	233

and the Fatty acid transporter 4	
5.2.8 Investigating the carbohydrate-binding properties of nine proteins selected from the <i>P. falciparum</i> merozoite surface protein library	233
5.3 Discussion	246
5.3.1 Investigating the binding of a library of <i>P. falciparum</i> merozoite surface proteins to human erythrocytes using a novel high-throughput, flow cytometry-based approach	247
5.3.2 Screening <i>P. falciparum</i> merozoite surface proteins against a panel of erythrocyte multi-pass receptors expressed recombinantly on HEK293E cells, to identify novel interactions	249
5.3.2.1 Endogenous receptors on HEK293E cells were recognised by some <i>P. falciparum</i> proteins	249
5.3.2.2 A number of potential erythrocyte multi-pass receptor: <i>P. falciparum</i> ligand pairs were identified in the screen	251
5.3.2.3 Further validation/characterisation of putative interactions between parasite proteins and recombinant multi-pass receptors.	252
5.3.3 The glycan-binding properties of a selected subset of merozoite surface proteins were tested by screening against a panel of synthetic carbohydrate probes.	253
5.4 Conclusion	254
5.5 Bibliography	255